




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C07D 239/42, 239/48, 401/04, 417/04, 409/04, A61K 31/505</p>	<p>A1</p>	<p>(11) International Publication Number: WO 97/19065 (43) International Publication Date: 29 May 1997 (29.05.97)</p>
<p>(21) International Application Number: PCT/GB96/02854 (22) International Filing Date: 20 November 1996 (20.11.96) (30) Priority Data: 9523675.8 20 November 1995 (20.11.95) GB (71) Applicant (for all designated States except US): CELLTECH THERAPEUTICS LIMITED [GB/GB]; 216 Bath Road, Slough, Berkshire SL1 4EN (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): DAVIS, Peter, David [GB/GB]; 10 Aston Park, Aston Rowant, Oxford OX9 5SW (GB). MOFFAT, David, Festus, Charles [GB/GB]; 14 Lonsdale Way, Holyport, Maidenhead, Berkshire SL6 2YX (GB). DAVIS, Jeremy, Martin [GB/GB]; 6 Marlborough Court, Wiltshire Drive, Wokingham, Berkshire RG40 1TA (GB). HUTCHINGS, Martin, Clive [GB/GB]; 93 Roycroft Lane, Barkham, Wokingham, Berkshire RG40 4NH (GB). (74) Agents: HALLYBONE, Huw, George et al.; Carpmals & Ransford, 43 Bloomsbury Square, London WCA 2RA (GB).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i></p>
<p>(54) Title: SUBSTITUTED 2-ANILINOPYRIMIDINES USEFUL AS PROTEIN KINASE INHIBITORS (57) Abstract Compounds of general formula (1) are described wherein R¹ is a hydrogen or halogen atom or an optionally substituted straight or branched chain alkyl, alkenyl or alkynyl group or a group selected from hydroxyl (-OH), substituted hydroxyl, thiol (-SH), substituted thiol, amino (-NH₂), or substituted amino; R² and R³, which may be the same or different, is each an optionally substituted straight or branched chain alkyl, alkenyl or alkynyl group; R⁴ is a hydrogen atom or a straight or branched chain alkyl group; R⁵ is a hydrogen atom or an optionally substituted straight or branched chain alkyl, alkenyl or alkynyl group; R⁶ is a hydrogen or halogen atom or an amino (-NH₂), substituted amino, nitro, carboxyl (-CO₂H) or esterified carboxyl group or a group -X¹-R^{6a} where X¹ is a direct bond or a linker atom or group and R^{6a} is an optionally substituted straight or branched chain alkyl, alkenyl or alkynyl group; X is a direct bond or a linker atom or group; R⁷ is an optionally substituted aliphatic, cycloaliphatic, heteroaliphatic, heterocycloaliphatic, aromatic or heteroaromatic group; and the salts, solvates, hydrates and N-oxides thereof. The compounds are selective protein kinase inhibitors, particularly the kinases p56^{lck}, p59^{fyn}, ZAP-70 and protein kinase C and are of use in the prophylaxis and treatment of immune diseases, hyperproliferative disorders and other diseases in which inappropriate protein kinase action is believed to have a role.</p> <div style="text-align: center;">  <p>(1)</p> </div>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

**SUBSTITUTED 2-ANILINOPYRIMIDINES USEFUL AS PROTEIN
KINASE INHIBITORS**

- 5 This invention relates to substituted 2-anilinopyrimidines, to processes for their preparation, to pharmaceutical compositions containing them, and to their use in medicine.

10 Protein kinases participate in the signalling events which control the activation, growth and differentiation of cells in response to extracellular mediators and to changes in the environment. In general, these kinases fall into two groups; those which preferentially phosphorylate serine and/or threonine residues and those which preferentially phosphorylate tyrosine residues [Hanks, S K, Hunter T, FASEB. J. 9, 576-596 (1995)].

15 The serine/threonine kinases include for example, protein kinase C isoforms [Newton A C, J. Biol. Chem. 270, 28495-28498 (1995)] and a group of cyclin-dependent kinases such as cdc2 [Pines J, Trends in Biochemical Sciences 18, 195-197 (1995)]. The tyrosine kinases include

20 membrane-spanning growth factor receptors such as the epidermal growth factor receptor [Iwashita S and Kobayashi M. Cellular Signalling 4, 123-132 (1992)], and cytosolic non-receptor kinases such as p56^{lck} p59^{fyn} ZAP-70 and csk kinases [Chan C *et al* Ann. Rev. Immunol. 12, 555-592 (1994)].

25 Inappropriately high protein kinase activity has been implicated in many diseases resulting from abnormal cellular function. This might arise either directly or indirectly, for example by failure of the proper control mechanisms for the kinase, related for example to mutation, overexpression or inappropriate activation of the enzyme; or by over- or


30 underproduction of cytokines or growth factors also participating in the transduction of signal upstream or downstream of the kinase. In all of these instances, selective inhibition of the action of the kinase might be expected to have a beneficial effect.

35 We have now found a series of substituted 2-anilinopyrimidines which are potent and selective inhibitors of protein kinases, especially the kinases

p56^{lck}, p59^{fyn}, ZAP-70 and protein kinase C. The compounds are thus of use in the prophylaxis and treatment of immune diseases, hyperproliferative disorders and other diseases in which inappropriate protein kinase action is believed to have a role.

5

Thus, according to one aspect of the invention, we provide a compound of formula (1):



10

wherein R¹ is a hydrogen or halogen atom or an optionally substituted straight or branched chain alkyl, alkenyl or alkynyl group or a group selected from hydroxyl (-OH), substituted hydroxyl, thiol (-SH), substituted thiol, amino (-NH₂), or substituted amino;

15 R² and R³, which may be the same or different, is each an optionally substituted straight or branched chain alkyl, alkenyl or alkynyl group;

R⁴ is a hydrogen atom or a straight or branched chain alkyl group;

R⁵ is a hydrogen atom or an optionally substituted straight or branched chain alkyl, alkenyl or alkynyl group;

20 R⁶ is a hydrogen or halogen atom or an amino (-NH₂), substituted amino, nitro, carboxyl (-CO₂H) or esterified carboxyl group or a group -X¹-R^{6a} where X¹ is a direct bond or a linker atom or group and R^{6a} is an optionally substituted straight or branched chain alkyl, alkenyl or alkynyl group;

25 X is a direct bond or a linker atom or group;

R⁷ is an optionally substituted aliphatic, cycloaliphatic, heteroaliphatic, heterocycloaliphatic, aromatic or heteroaromatic group; and the salts, solvates, hydrates and N-oxides thereof.

Halogen atoms represented by the group R^1 and/or R^6 in compounds of formula (1) include for example fluorine, chlorine, bromine or iodine atoms.

- 5 When R^1 , R^2 , R^3 , R^5 and/or R^{6a} is an optionally substituted straight or branched chain alkyl, alkenyl or alkynyl group each of said groups may independently be an optionally substituted straight or branched chain C_{1-6} alkyl, e.g. C_{1-3} alkyl, C_{2-6} alkenyl, e.g. C_{2-4} alkenyl, or C_{2-6} alkynyl, e.g. C_{2-4} alkynyl group. Particular examples of such groups include optionally substituted $-CH_3$, $-CH_2CH_3$, $-(CH_2)_2CH_3$, $-CH(CH_3)_2$, $-(CH_2)_3CH_3$, $-CH(CH_3)CH_2CH_3$, $-CH_2CH(CH_3)_2$, $-C(CH_3)_3$, $-(CH_2)_4CH_3$, $-(CH_2)_5CH_3$, $-CHCH_2$, $-CHCHCH_3$, $-CH_2CHCH_2$, $-CHCHCH_2CH_3$, $-CH_2CHCHCH_3$, $-(CH_2)_2CHCH_2$, $-CCH$, $-CCCH_3$, $-CH_2CCH$, $-CCCH_2CH_3$, $-CH_2CCCH_3$, or $-(CH_2)_2CCH$ groups. The optional substituents which may be present on these groups include one, two, three or more substituents selected from halogen atoms, e.g. fluorine, chlorine, bromine or iodine atoms, or hydroxyl, C_{1-6} alkoxy, e.g. methoxy or ethoxy, thiol, C_{1-6} alkylthio e.g. methylthio or ethylthio, amino, C_{1-6} alkylamino, e.g. methylamino or ethylamino, or C_{1-6} dialkylamino, e.g. dimethylamino or diethylamino groups.

- Substituted hydroxyl groups represented by the group R^1 in compounds of formula (1) include $-OR^8$ groups where R^8 is an optionally substituted straight or branched C_{1-6} alkyl, e.g. methyl or ethyl, C_{2-6} alkenyl, e.g. allyl, or C_{2-6} alkynyl, e.g. ethynyl group. The optional substituents which may be present on these groups include a phenyl group and/or one, two, three or more of the atoms or groups described above in relation to substituents present on alkyl groups represented by R^1 .

- 30 Substituted thiol groups represented by the group R^1 include $-SR^8$ groups, wherein R^8 is as just defined.

- When R^1 and/or R^6 is a substituted amino group it may be for example a $-NHR^9$ or $-NR^9R^{10}$ group where R^9 and R^{10} , which may be the same or different, is each a group $-R^8$ or $-COR^8$ where R^8 is as just defined.

Esterified carboxyl groups represented by the group R⁶ include groups of formula -CO₂Alk¹ wherein -CO₂Alk¹ is as defined hereinafter in connection with esterified carboxyl groups represented by the group R¹³.

- 5 Linker atoms represented by X or X¹ in compounds of formula (1) include -O- or -S- atoms. When X or X¹ is a linker group it may be for example a -C(O)-, -C(S)-, -S(O)-, -S(O)₂-, -N(R¹¹)- [where R¹¹ is a hydrogen atom or a C₁₋₆ alkyl, e.g. methyl or ethyl, group], -CON(R¹¹)-, -OC(O)N(R¹¹)-, -CSN(R¹¹)-, -N(R¹¹)CO-, -N(R¹¹)C(O)O-, -N(R¹¹)CS-, -SON(R¹¹),
10 -SO₂N(R¹¹), -N(R¹¹)SO₂-, -N(R¹¹)CON(R¹¹)-, -N(R¹¹)CSN(R¹¹)-, -N(R¹¹)SON(R¹¹)- or -N(R¹¹)SO₂N(R¹¹) group.

- When R⁷ in compounds of formula (1) is an optionally substituted aliphatic or cycloaliphatic group it may be an optionally substituted C₁₋₁₀ aliphatic
15 or C₃₋₁₀ cycloaliphatic group. Particular examples include optionally substituted straight or branched chain C₁₋₆ alkyl, C₂₋₆ alkenyl, or C₂₋₆ alkynyl groups or optionally substituted C₃₋₁₀cycloalkyl, C₃₋₁₀cycloalkenyl or C₃₋₁₀cycloalkynyl groups.

- 20 Heteroaliphatic or heterocycloaliphatic groups represented by R⁷ include the aliphatic or cycloaliphatic groups just described but with each group additionally containing one, two, three or four heteroatoms or heteroatom-containing groups. Particular heteroatoms or groups include atoms or groups -X²- where X² is as defined above for X when X is a linker atom or
25 group.

- Aromatic groups represented by R⁷ in compounds of formula (1) include for example optionally substituted monocyclic or bicyclic fused ring C₆₋₁₂ aromatic groups, such as optionally substituted phenyl, 1- or 2-naphthyl, 1-
30 or 2-tetrahydronaphthyl, indanyl or indenyl groups.

- Heteroaromatic groups represented by R⁷ include optionally substituted C₁₋₉ heteroaromatic groups containing for example one, two, three or four heteroatoms selected from oxygen, sulphur or nitrogen atoms. In general,
35 the heteroaromatic groups may be for example monocyclic or bicyclic fused ring heteroaromatic groups. Monocyclic heteroaromatic groups

include for example five- or six-membered heteroaromatic groups containing one, two, three or four heteroatoms selected from oxygen, sulphur or nitrogen atoms. Bicyclic heteroaromatic groups include for example nine- to thirteen-membered fused-ring heteroaromatic groups
5 containing one, two or more heteroatoms selected from oxygen, sulphur or nitrogen atoms.

In general, when R^7 is a heteroaliphatic, heterocycloaliphatic or heteroaromatic group it is attached to the remainder of the molecule of
10 formula (1) through any available heteroatom or group or, preferably, carbon atom.

Particular examples of R^7 aliphatic groups include those alkyl, alkenyl or alkynyl groups specifically described above in relation to the groups R^1 ,
15 R^2 , R^3 , R^5 and R^6 . Each of these groups may be optionally substituted, and/or optionally interrupted by one or two heteroatoms or heteroatom-containing groups represented by $-X^2-$ [where X^2 is as previously defined], to yield particular examples of R^7 optionally substituted aliphatic or heteroaliphatic groups.

20 Particular examples of R^7 cycloaliphatic and heterocycloaliphatic groups include optionally substituted cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, 2-cyclobuten-1-yl, 2-cyclopenten-1-yl, 3-cyclopenten-1-yl, 2,4-cyclopentadien-1-yl, 3,5,-cyclohexadien-1-yl, pyrroline, e.g. 2- or 3-pyrrolinyl, pyrrolidinyl, dioxolanyl, e.g. 1,3-dioxolanyl, imidazoliny, e.g. 2-imidazoliny, imidazolidiny, pyrazoliny, e.g. 2-pyrazoliny, pyrazolidiny, pyranyl, e.g. 2- or 4-pyranyl, piperidinyl, 1,4-dioxanyl, morpholiny, 1,4-dithianyl, thiomorpholiny, piperazinyl, 1,3,5-trithianyl, oxaziny, e.g. 2H-1,3-, 6H-1,3-, 6H-1,2-, 2H-1,2- or 4H-1,4-oxaziny, 1,2,5-oxathiaziny, isoxaziny, e.g. o- or p-isoxaziny, oxathiaziny, e.g. 1,2,5 or 1,2,6-oxathiaziny, or 1,3,5,2-oxadiaziny groups.

Examples of heteroaromatic groups represented by R^7 include optionally substituted pyrrolyl, furyl, thienyl, imidazolyl, N-methylimidazolyl, N-ethyl-
35 imidazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyrazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl,

1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, 1,3,4-thiadiazole, pyridyl, pyrimidinyl, pyridazinyl, pyrazinyl, 1,3,5-triazinyl, 1,2,4-triazinyl, 1,2,3-triazinyl, benzofuryl, [2,3-dihydro]benzofuryl, isobenzofuryl, benzothienyl, benzotriazolyl, isobenzothienyl, indolyl, isoindolyl, benzimidazolyl, imidazo[1,2-a]pyridyl, benzothiazolyl, benzoxazolyl, benzopyranyl, [3,4-dihydro]benzopyranyl, quinazolinyl, naphthyridinyl, pyrido[3,4-b]pyridyl, pyrido[3,2-b]pyridyl, pyrido[4,3-b]pyridyl, quinolinyl, isoquinolinyl, tetrazolyl, 5,6,7,8-tetrahydroquinolinyl, 5,6,7,8-tetrahydroisoquinolinyl, and imidyl, e.g. succinimidyl, phthalimidyl, or naphthalimidyl such as 1,8-naphthalimidyl.

Optional substituents which may be present on any of the above R⁷ groups in compounds of formula (1) include one, two, three or more substituents, each represented by the group R¹². The substituent R¹² may be selected from an atom or group R¹³ or -Alk(R¹³)_m, where R¹³ is a halogen atom, or an amino (-NH₂), -NHR¹⁴ [where R¹⁴ is an -Alk(R¹³)_m, heterocycloalkyl, -Alk-heterocycloalkyl, aryl or heteroaryl group], -N(R¹⁴)₂ [where each R¹⁴ group is the same or different], nitro, cyano, hydroxyl (-OH), -OR¹⁴, formyl, carboxyl (-CO₂H), esterified carboxyl, thiol (-SH), -SR¹⁴, -COR¹⁴, -CSR¹⁴, -SO₃H, -SO₂R¹⁴, -SO₂NH₂, -SO₂NHR¹⁴, SO₂N[R¹⁴]₂, -CONH₂, -CSNH₂, -CONHR¹⁴, -CSNHR¹⁴, -CON[R¹⁴]₂, -CSN[R¹⁴]₂, -NHSO₂H, -NHSO₂R¹⁴, -N[SO₂R¹⁴]₂, -NHSO₂NH₂, -NHSO₂NHR¹⁴, -NHSO₂N[R¹⁴]₂, -NHCOR¹⁴, -NHCONH₂, -NHCONHR¹⁴, -NHCON[R¹⁴]₂, -NHCSR¹⁴, -NHCSNH₂, -NHCSNHR¹⁴, -NHCSN[R¹⁴]₂, -NHC(O)OR¹⁴, or an optionally substituted cycloalkyl aryl or heteroaryl group; or R¹² may be an optionally substituted heterocycloalkyl or -Alk-heterocycloalkyl group provided that when R¹² is a heterocycloalkyl group X in compounds of the invention is either a linker atom or group, or X is a bond and R⁷ is an optionally substituted aliphatic, cycloaliphatic, heteroaliphatic, heterocycloaliphatic, aromatic or heteroaromatic group other than a pyridyl group; Alk is a straight or branched C₁₋₆ alkylene, C₂₋₆ alkenylene or C₂₋₆ alkynylene chain, optionally interrupted by one, two or three -O- or -S- atoms or S-(O)-, -S(O)₂- or -N(R¹¹)- groups; and m is zero or an integer 1, 2 or 3.

35

When in the group $-\text{Alk}(\text{R}^{13})_m$ m is an integer 1, 2 or 3, it is to be understood that the substituent or substituents R^{13} may be present on any suitable carbon atom in $-\text{Alk}$. Where more than one R^{13} substituent is present these may be the same or different and may be present on the same or different atom in $-\text{Alk}$ or in R^7 as appropriate. Thus for example, R^7 may represent a $-\text{CH}(\text{R}^{13})_2$ group, such as a $-\text{CH}(\text{OH})\text{Ar}$ group where Ar is an aryl or heteroaryl group as defined below. Clearly, when m is zero and no substituent R^{13} is present the alkylene, alkenylene or alkynylene chain represented by Alk becomes an alkyl, alkenyl or alkynyl group.

10

When R^{13} is a halogen atom it may be for example a fluorine, chlorine, bromine, or iodine atom.

Esterified carboxyl groups represented by the group R^{13} include groups of formula $-\text{CO}_2\text{Alk}^1$ wherein Alk^1 is a straight or branched, optionally substituted C_{1-8} alkyl group such as a methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl or t-butyl group; a $\text{C}_{6-12}\text{arylC}_{1-8}\text{alkyl}$ group such as an optionally substituted benzyl, phenylethyl, phenylpropyl, 1-naphthylmethyl or 2-naphthylmethyl group; a $\text{C}_{6-12}\text{aryl}$ group such as an optionally substituted phenyl, 1-naphthyl or 2-naphthyl group; a $\text{C}_{6-12}\text{aryloxyC}_{1-8}\text{alkyl}$ group such as an optionally substituted phenyloxymethyl, phenyloxyethyl, 1-naphthyloxymethyl, or 2-naphthyloxymethyl group; an optionally substituted $\text{C}_{1-8}\text{alkanoyloxyC}_{1-8}\text{alkyl}$ group, such as a pivaloyloxymethyl, propionyloxyethyl or propionyloxypropyl group; or a $\text{C}_{6-12}\text{aroyloxyC}_{1-8}\text{alkyl}$ group such as an optionally substituted benzoyloxyethyl or benzoyloxypropyl group. Optional substituents present on the Alk^1 group include R^{13} substituents described above.

When Alk is present in or as a substituent R^{12} it may be for example a methylene, ethylene, n-propylene, i-propylene, n-butylene, i-butylene, s-butylene, t-butylene, ethenylene, 2-propenylene, 2-butenylene, 3-butenylene, ethynylene, 2-propynylene, 2-butynylene or 3-butynylene chain, optionally interrupted by one, two, or three $-\text{O}-$ or $-\text{S}-$, atoms or $-\text{S}(\text{O})-$, $-\text{S}(\text{O})_2-$ or $-\text{N}(\text{R}^{11})-$ groups.

35

Optionally substituted cycloalkyl groups represented by the group R¹³ include optionally substituted C₅₋₇ cycloalkyl groups such as optionally substituted cyclopentyl or cyclohexyl groups.

- 5 Heterocycloalkyl groups represented by the group R¹² or R¹⁴ include optionally substituted heteroC₃₋₆cycloalkyl groups containing one or two oxygen, sulphur or nitrogen atoms. Particular examples of such groups include optionally substituted azetidiny, pyrrolidiny, piperidiny, piperaziny, homopiperaziny, morpholiny or thiomorpholiny groups. The heterocyclo-
- 10 alkyl group may be attached to the remainder of the molecule through any of its ring carbon atoms, or where present, ring nitrogen atom. Where the group R¹² is an -Alk-heterocycloalkyl group, Alk may be as defined above and the heterocycloalkyl portion may be as just defined, attached to Alk through any of its ring carbon atoms, or where present, ring nitrogen atom.

- 15 Optional substituents which may be present on R¹², R¹³ or R¹⁴ cycloalkyl or heterocycloalkyl groups include one or two C₁₋₆ alkyl, e.g. methyl or ethyl, hydroxyl (-OH) hydroxyC₁₋₆alkyl, e.g. hydroxymethyl or hydroxyethyl, or C₁₋₆ alkoxy, e.g. methoxy or ethoxy groups. The substituent(s) may be
- 20 present on any available ring carbon or nitrogen atom as appropriate.

- Aryl and heteroaryl groups represented by the groups R¹³ or R¹⁴ include for example optionally substituted monocyclic or bicyclic C₆₋₁₂ aromatic groups, or C₁₋₉ heteroaromatic groups such as those described above in
- 25 relation to the group R⁷.

- Particularly useful atoms or groups represented by R¹² include fluorine, chlorine, bromine or iodine atoms, or C₁₋₆alkyl, C₁₋₆ alkylamino, C₁₋₆hydroxyalkyl, C₁₋₆alkylthiol, C₁₋₆alkoxy, hydroxyC₁₋₆alkoxy, aminoC₁₋₆alkoxy, C₁₋₆alkylaminoC₁₋₆alkoxy, C₁₋₆dialkylaminoC₁₋₆alkoxy, optionally
- 30 substituted C₅₋₇cyclo-alkoxy, optionally substituted C₅₋₇cycloalkyl, optionally substituted C₅₋₇cycloalkylamino, haloC₁₋₆alkyl, haloC₁₋₆alkoxy, C₁₋₆alkylamino, amino (-NH₂), aminoC₁₋₆alkyl, C₁₋₆dialkylamino, hydroxyC₁₋₆ alkylamino, aminoC₁₋₆alkylamino, C₁₋₆alkylaminoC₁₋₆alkylamino, C₁₋₆dialkylaminoC₁₋₆alkylamino, C₁₋₆alkylaminoC₁₋₆
- 35 dialkylamino, C₁₋₆dialkylaminoC₁₋₆dialkylamino, nitro, cyano, hydroxyl (-

OH), formyl [HC(O)-], carboxyl (-CO₂H), -CH₂CO₂H, -OCH₂CO₂H, -CO₂Alk¹ [where Alk¹ is as defined above], -CH₂CO₂Alk¹, C₁₋₆alkoxycarbonylC₁₋₆alkoxy, C₁₋₆alkanoyl, optionally substituted phenyl C₁₋₆alkanoyl, thiol (-SH), thioC₁₋₆alkyl, -SC(NH)NH₂, sulphonyl (-SO₃H), C₁₋₆alkylsulphonyl, optionally substituted phenylsulphonyl, aminosulphonyl (-SO₂NH₂), C₁₋₆alkylaminosulphonyl, C₁₋₆dialkylaminosulphonyl, optionally substituted phenylamino-sulphonyl, carboxamido (-CONH₂), C₁₋₆alkylaminocarbonyl, C₁₋₆dialkylaminocarbonyl, optionally substituted phenylaminocarbonyl, aminocarbonylmethyl, C₁₋₆alkylaminocarbonylmethyl, optionally substituted benzylaminocarbonylmethyl, -NHC(S)NH₂, sulphonylamino (-NH₂SO₂H), C₁₋₆alkylsulphonylamino, C₁₋₆dialkylsulphonylamino, optionally substituted phenylsulphonylamino, amino-sulphonylamino (-NH₂SO₂NH₂), C₁₋₆alkylaminosulphonylamino, C₁₋₆dialkylaminosulphonylamino, optionally substituted phenylaminosulphonylamino, aminocarbonylamino, C₁₋₆alkylaminocarbonylamino, C₁₋₆dialkylaminocarbonylamino, phenylaminocarbonylamino, C₁₋₆alkanoylamino, aminoC₁₋₆alkanoylamino, optionally substituted pyridylcarboxyamino, C₁₋₆alkanoylaminoC₁₋₆alkyl, C₁₋₆alkoxycarbonylamino, optionally substituted heteroC₃₋₆cycloalkyl, piperidiny, piperaziny, 4-methyl-piperaziny, homopiperaziny, or morpholiny, optionally substituted heteroC₃₋₆cycloalkylC₁₋₆alkyl, piperidinyC₁₋₆alkyl, piperazinyC₁₋₆alkyl or morpholinyC₁₋₆alkyl, optionally substituted heteroC₃₋₆alkylC₁₋₆alkylamino, optionally substituted heteroC₃₋₆cycloalkylamino, tetrazoly, optionally substituted phenylamino, optionally substituted benzylamino, optionally substituted benzyloxy, or optionally substituted pyridylmethylamino group.

Where desired, two R¹² substituents may be linked together to form a cyclic group such as a cyclic ether, e.g. a C₁₋₆alkylenedioxy group such as a methylenedioxy or ethylenedioxy group.

30

Especially useful R¹² substituents include for example fluorine, chlorine, bromine or iodine atoms, or a methylamino, ethylamino, hydroxymethyl, hydroxyethyl, methylthiol, ethylthiol, methoxy, ethoxy, n-propoxy, 2-hydroxyethoxy, 3-hydroxypropoxy, 4-hydroxybutoxy, 2-aminoethoxy, 3-aminopropoxy, 2-(methylamino)ethoxy, 2-(dimethylamino)ethoxy, 3-(dimethylamino)propoxy, cyclopentyloxy, cyclohexyl, cyclohexylamino, 2-

35

hydroxycyclohexylamino, trifluoromethyl, trifluoromethoxy, methylamino, ethylamino, amino $(-\text{NH})_2$, aminomethyl, aminoethyl, dimethylamino, diethylamino, ethyl(methyl)amino, propyl(methyl)amino, 2-hydroxyethylamino, 3-hydroxypropylamino, 4-hydroxybutylamino, 2-aminoethylamino, 3-aminopropylamino, 4-aminobutylamino, 2-(methylamino)ethylamino, 2-(ethylamino)ethylamino, 2-(i-propylamino)ethylamino, 3-(i-propylamino)propylamino, 2-(dimethylamino)ethylamino, 3-(dimethylamino)propylamino, 2-(diethylamino)ethylamino, 3-(diethylamino)propylamino, 2-(methylamino)-ethyl(methyl)amino, 3-(methylamino)propyl(methyl)amino, 2-(dimethylamino)ethyl(methyl)amino, 2-(dimethylamino)ethyl(ethyl)amino, nitro, cyano, hydroxyl $(-\text{OH})$, formyl $[\text{HC}(\text{O})-]$, carboxyl $(-\text{CO}_2\text{H})$, $-\text{CH}_2\text{CO}_2\text{H}$, $-\text{OCH}_2\text{CO}_2\text{H}$, $-\text{CO}_2\text{CH}_3$, $-\text{CO}_2\text{CH}_2\text{CH}_3$, $-\text{CH}_2\text{CO}_2\text{CH}_3$, $-\text{CH}_2\text{CO}_2\text{CH}_2\text{CH}_3$, $-\text{CH}_2\text{CO}_2\text{CH}_2\text{phenyl}$, t-butoxycarbonylmethoxy, acetyl, phenacetyl, thio $(-\text{SH})$, thiomethyl, thioethyl, $-\text{SC}(\text{NH})\text{NH}_2$, sulphonyl $(-\text{SO}_2\text{H})$, methylsulphonyl, methylaminosulphonyl, ethylaminosulphonyl, dimethylaminosulphonyl, diethylaminosulphonyl, carboxamido $(-\text{CONH}_2)$, methylaminocarbonyl, ethylaminocarbonyl, dimethylaminocarbonyl, diethylaminocarbonyl, methylaminocarbonylmethyl, $-\text{NHC}(\text{S})\text{NH}_2$, sulphonylamino $(-\text{NHSO}_2\text{H})$, methylsulphonylamino, ethylsulphonylamino, dimethylsulphonylamino, diethylsulphonylamino, sulphonylamino $(-\text{HHSO}_2\text{NH}_2)$, methylaminosulphonylamino, ethylaminosulphonylamino, dimethylaminosulphonylamino, diethylaminosulphonylamino, methylaminocarbonylamino, ethylaminocarbonylamino, dimethylaminocarbonylamino, diethylaminocarbonylamino, acetylamino, aminomethylcarbonylamino, acetylamino-methyl, methoxycarbonylamino, ethoxycarbonylamino, t-butoxycarbonylamino, pyrrolidinyl, piperidinyl, piperazinyl, 4-methylpiperazinyl, homopiperazinyl, morpholinyl, pyrrolidinyl $\text{C}_{1-6}\text{alkyl}$, piperidinyl $\text{C}_{1-6}\text{alkyl}$, piperazinyl $\text{C}_{1-6}\text{alkyl}$, morpholinyl $\text{C}_{1-6}\text{alkyl}$, 2-pyrrolidinyethylamino, 2-(1-methylpyrrolidinyl)ethylamino, 1-ethylpyrrolidinylmethylamino, piperidinylamino, 1-benzylpiperidinylamino, 4-(methoxy)phenylamino, 4-(3-hydroxypropyl)phenylamino, benzylamino, benzyloxy, pyridylmethylamino group.

It will be appreciated that where two or more R^{12} substituents are present, these need not necessarily be the same atoms and/or groups.

The presence of certain substituents in the compounds of formula (1) may enable salts of the compounds to be formed. Suitable salts include pharmaceutically acceptable salts, for example acid addition salts derived from inorganic or organic acids, and salts derived from inorganic and organic bases.

Acid addition salts include hydrochlorides, hydrobromides, hydroiodides, alkylsulphonates, e.g. methanesulphonates, ethanesulphonates, or isethionates, arylsulphonates, e.g. p-toluenesulphonates, besylates or napsylates, phosphates, sulphates, hydrogen sulphates, acetates, trifluoroacetates, propionates, citrates, maleates, fumarates, malonates, succinates, lactates, oxalates, tartrates and benzoates.

Salts derived from inorganic or organic bases include alkali metal salts such as sodium or potassium salts, alkaline earth metal salts such as magnesium or calcium salts, and organic amine salts such as morpholine, piperidine, dimethylamine or diethylamine salts.

Particularly useful salts of compounds according to the invention include pharmaceutically acceptable salts, especially acid addition pharmaceutically acceptable salts.

It will be appreciated that depending on the nature of the substituents R¹-R³ and R⁵ - R⁷ the compounds of formula (1) may exist as geometrical isomers and/or may have one or more chiral centres so that enantiomers or diastereomers may exist. It is to be understood that the invention extends to all such isomers of the compounds of formula (1), and to mixtures thereof, including racemates.

In one class of compounds of formula (1) the groups R¹, R², R³, R⁴, R⁵, R⁷ and X are as defined for formula (1), as is the group R⁶ except it is not an amino, substituted amino, nitro, carboxyl or esterified carboxyl group.

One particular class of compounds according to the invention has the formula (1) wherein R² and R³ is each an optionally substituted methyl or ethyl group. Particular examples of groups of these types include methyl,

ethyl, halomethyl, e.g. $-\text{CH}_2\text{F}$, $-\text{CH}_2\text{Cl}$, $-\text{CHF}_2$, $-\text{CHCl}_2$, $-\text{CF}_3$, $-\text{CCl}_3$, or haloethyl groups. In general however, R^2 and R^3 is each preferably a methyl group. In compounds of this class and in general in compounds of formula (1) R^1 may in particular be an optionally substituted methoxy group. Particular examples of such R^1 groups include $-\text{OCH}_2\text{F}$, $-\text{OCH}_2\text{Cl}$, $-\text{OCHF}_2$, $-\text{OCHCl}_2$, $-\text{OCF}_3$, $-\text{OCCl}_3$ or, especially, methoxy groups.


In a further preferred class of compounds of formula (1) R^4 is preferably a hydrogen atom.

The groups R^5 and R^6 in compounds of formula (1) are each preferably a hydrogen atom.

In yet another preference, X in compounds of formula (1) is a direct bond, an oxygen or sulphur atom or a $-\text{N}(\text{R}^{11})-$ group. Especially useful compounds of this type are those wherein X is a direct bond, a sulphur atom or a $-\text{N}(\text{R}^{11})-$, particularly a $-\text{NH}-$, group.

R^7 in compounds of formula (1) is preferably an optionally substituted aromatic or heteroaromatic group.

A further class of compounds according to the invention has the formula 1(a):



1(a)

wherein R^5 and R^6 are as defined for formula (1), X is a direct bond, an oxygen or sulphur atom, or a group $-\text{N}(\text{R}^{11})-$ and R^7 is an optionally

substituted aromatic or heteroaromatic group, and the salts, solvates, hydrates and N-oxides thereof.

5 In these compounds, R⁵ is preferably a hydrogen atom. R⁶ is preferably a group -X¹R^{6a} where X¹ is as defined for formula (1) and R^{6a} is an optionally substituted straight or branched chain alkyl group, or R⁶ is especially a hydrogen atom.

10 X in compounds of formula (1a) is preferably a direct bond, a sulphur atom, or a -N(R¹¹)- group, particularly a -NH-group.

15 The aromatic or heteroaromatic R⁷ group in compounds of formulae (1) or 1(a) in general may be as defined previously for compounds of formula (1). In one preference, however, R⁷ is an optionally substituted phenyl, 1- or 2-naphthyl or heteroaromatic group containing one or two oxygen, sulphur and/or nitrogen atoms. Thus in particular R⁷ may be an optionally substituted phenyl, 1- or 2-naphthyl, pyrrolyl, furyl, thienyl, indolyl, pyrazolyl, thiazolyl, [2,3-dihydro]benzofuryl, benzothiazolyl, 2-pyridyl, 3-pyridyl or 4-pyridyl group. Particularly useful groups include optionally substituted phenyl, 2-pyridyl, 3-pyridyl or 4-pyridyl groups. The aromatic or heteroaromatic group may in particular be attached to the remainder of the compound of formula (1) through any available ring carbon atom.

25 In general, the optional substituents which may be present on aromatic or heteroaromatic R⁷ groups in compounds of formulae (1) or (1a) include one, two, or three R¹² substituents as generally and particularly described above and hereinafter in the Examples. Particularly useful R¹² substituents include -NHR¹⁴, -AlkNH₂, -AlkNHR¹⁴, -OR¹⁴, -AlkCO₂H or -AlkCO₂Alk¹ groups where R¹⁴, Alk and Alk¹ are as generally and particularly defined above. Useful members of these substituents include those wherein R¹⁴ is an -Alk, -AlkNH₂ or -Alk-heterocycloalkyl group. In these, and the other preferred substituents just mentioned, Alk and Alk¹ when present is each preferably a C₁₋₆alkyl group.

35 Particularly useful compounds according to the invention include:
N,N'-Bis(3,4,5-Trimethoxyphenyl)-2,4-pyrimidinediamine;

- 4-(2-(2-Dimethylaminoethylamino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine;
- 4-(2-(4-Hydroxybutylamino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine;
- 5 4-(3-(3-Aminopropoxy)phenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine;
- N4-(3,4-Dimethoxyphenyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine;
- 4-(4-(2-Hydroxyethoxy)phenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine;
- 10 4-(2-(3-(Morpholino)propylamino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine;
- 4-(2-(2-(1-Methylpyrrolidin-2-yl)ethylamino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine;
- 15 N4-(4-(Ethoxycarbonylmethyl)phenyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine;
- N4-(4-Hydroxyphenyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine and
- N4-(4-(3-Aminopropoxy)phenyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine;
- 20 and the salts, solvates, hydrates and N-oxides thereof

Compounds according to the invention are potent and selective inhibitors of protein kinases as demonstrated by differential inhibition of enzymes

25 such as EGFr kinase, p56^{lck} kinase, ZAP-70 kinase, Csk kinase and p59^{fyn} kinase. The ability of the compounds to act in this way may be simply determined by employing tests such as those described in the Examples hereinafter.

30 The compounds according to the invention are thus of particular use in the prophylaxis and treatment of diseases in which inappropriate protein tyrosine kinase action plays a role, for example in autoimmune diseases such as rheumatoid arthritis, multiple sclerosis, and systemic lupus erythematosus, in transplant rejection, in graft v host disease, in

35 hyperproliferative disorders such as tumours, psoriasis, in pannus formation in rheumatoid arthritis, restenosis following angioplasty and

atherosclerosis, and in diseases in which cells receive pro-inflammatory signals such as asthma, inflammatory bowel disease and pancreatitis.

5 For the prophylaxis or treatment of disease the compounds according to the invention may be administered as pharmaceutical compositions, and according to a further aspect of the invention we provide a pharmaceutical composition which comprises a compound of formula (1) together with one or more pharmaceutically acceptable carriers, excipients or diluents.

10 Pharmaceutical compositions according to the invention may take a form suitable for oral, buccal, parenteral, nasal, topical or rectal administration, or a form suitable for administration by inhalation or insufflation.

15 For oral administration, the pharmaceutical compositions may take the form of, for example, tablets, lozenges or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (e.g. pregelatinised maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose); fillers (e.g. lactose, microcrystalline cellulose or calcium hydrogen phosphate); lubricants (e.g. magnesium stearate, talc or silica); disintegrants (e.g. potato starch or sodium glycollate); or wetting agents (e.g. sodium lauryl sulphate). The tablets
20 may be coated by methods well known in the art. Liquid preparations for oral administration may take the form of, for example, solutions, syrups or suspensions, or they may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations
25 may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents, emulsifying agents, non-aqueous vehicles and preservatives. The preparations may also contain buffer salts, flavouring, colouring and sweetening agents as appropriate.

30 Preparations for oral administration may be suitably formulated to give controlled release of the active compound.

35 For buccal administration the compositions may take the form of tablets or lozenges formulated in conventional manner.

The compounds for formula (1) may be formulated for parenteral administration by injection e.g. by bolus injection or infusion. Formulations for injection may be presented in unit dosage form, e.g. in glass ampoule or multi dose containers, e.g. glass vials. The compositions for injection
5 may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilising, preserving and/or dispersing agents. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g. sterile pyrogen-free water, before use.

10 In addition to the formulations described above, the compounds of formula (1) may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation or by intramuscular injection.


15 For nasal administration or administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation for pressurised packs or a nebuliser, with the use of suitable propellant, e.g. dichlorodifluoromethane, trichloro-
20 fluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas or mixture of gases.

The compositions may, if desired, be presented in a pack or dispenser device which may contain one or more unit dosage forms containing the
25 active ingredient. The pack or dispensing device may be accompanied by instructions for administration.


The quantity of a compound of the invention required for the prophylaxis or treatment of a particular condition will vary depending on the compound
30 chosen, and the condition of the patient to be treated. In general, however, daily dosages may range from around 100ng/kg to 100mg/kg e.g. around 0.01mg/kg to 40mg/kg body weight for oral or buccal administration, from around 10ng/kg to 50mg/kg body weight for parenteral administration and around 0.05mg to around 1000mg e.g.
35 around 0.5mg to around 1000mg for nasal administration or administration by inhalation or insufflation.

The compounds of the invention may be prepared by a number of processes as generally described below and more specifically in the Examples hereinafter. In the following process description, the symbols R^1 - R^7 and X when used in the formulae depicted are to be understood to represent those groups described above in relation to formula (1) unless otherwise indicated. In the reactions described below, it may be necessary to protect reactive functional groups, for example hydroxy, amino, thio or carboxy groups, where these are desired in the final product, to avoid their unwanted participation in the reactions. Conventional protecting groups may be used in accordance with standard practice [see, for example, Green, T. W. in "Protective Groups in Organic Synthesis", John Wiley and Sons, 1981]. In some instances, deprotection may be the final step in the synthesis of a compound of formula (1) and the processes according to the invention described hereinafter are to be understood to extend to such removal of protecting groups.

Thus according to a further aspect of the invention, a compound of formula (1) wherein X is a direct bond may be prepared by reaction of a guanidine of formula (2):



or a salt thereof
with an enaminone of formula (3):



where R^{15} and R^{16} , which may be the same or different is each a C_{1-6} alkyl group.


The reaction may be performed in a solvent, for example a protic solvent such as an alcohol, e.g. ethanol, methoxyethanol or propanol, optionally in

the presence of a base e.g. an alkali metal base, such as sodium hydroxide or potassium carbonate, at an elevated temperature, e.g. the reflux temperature.

- 5 Salts of the compounds of formula (2) include acid salts such as inorganic acid salts e.g. hydrochlorides or nitrates.

Intermediate guanidines of formula (2) may be prepared by reaction of the corresponding aniline of formula (4):

10



with cyanamide at an elevated temperature.

- 15 The reaction may be performed in a solvent such as ethanol at an elevated temperature, e.g. up to the reflux temperature. Where it is desired to obtain a salt of a guanidine of formula (2), the reaction may be performed in the presence of a concentrated acid, e.g. hydrochloric or nitric acid.

20

The anilines of formula (4) are either known compounds or may be obtained by conventional procedures, for example by hydrogenation of the corresponding nitro derivatives using for example hydrogen in the presence of a metal catalyst in a suitable solvent, for example as more particularly described in the interconversion reactions discussed below or by use of the corresponding nitro derivative and a reducing agent such as a sodium hydrosulphite in a solvent such as ethanol at an elevated temperature such as the reflux temperature. The nitrobenzenes for this particular reaction are either known compounds or may be prepared using


25 similar methods to those used for the preparation of the known compounds, for example by treatment of the corresponding benzene with

30

nitric acid in the presence of an acid such as acetic acid at around ambient to the reflux temperature.

Intermediate enamminones of formula (3) may be prepared by reaction of an acetyl derivative $R^7COCH_2R^6$ with an acetal $(R^{16})(R^{15})NCR^5(OCH_3)_2$ at an elevated temperature. The starting materials for this reaction are either known compounds or may be prepared by methods analogous to those used for the preparation of the known compounds.

In another process according to the invention, a compound of formula (1) where X is a linker atom or group may be prepared by reaction of an intermediate of formula (5)



(5)

15

where L is a leaving atom or group, with a reagent R^7X^2H where X^2 is a linking atom or group as defined above.

Particular leaving atoms or groups represented by L include for example halogen atoms, e.g. bromine, iodine or chlorine atoms, and sulphonyloxy groups, e.g. alkylsulphonyloxy groups, such as trifluoromethylsulphonyloxy, and arylsulphonyloxy groups, such as p-toluenesulphonyloxy.

The reaction may be performed in the presence of a base, for example an organic base such as an organic amine, e.g. triethylamine or an inorganic base, for example a hydride such as sodium hydride, an alkoxide such as potassium t-butoxide, or a carbonate such as caesium or potassium carbonate, where necessary in the presence of a dipolar aprotic solvent

25


such as an ether, e.g. a cyclic ether such as tetrahydrofuran, or an amide, e.g. a substituted amide such as dimethylformamide, at a suitable temperature e.g. from around room temperature to around 90°C.

- 5 Intermediates of formula (5) may be prepared by reaction of the corresponding pyrimidinones (where $-N=C(L)-$ in formula (5) is $-NH-C(O)-$) with a halide using standard procedures, for example by reaction with a reagent $PO(Hal)_3$ (where Hal is a halogen atom), e.g. $POCl_3$, or RSO_2Hal (where R is an optionally substituted alkyl or aryl group) where necessary
10 at an elevated temperature.

The starting pyrimidinones may be prepared by reaction of a guanidine of formula (2) or a salt thereof with a β -ketoester [for example a compound $CH_3CH_2C(O)OCH(R^6)C(O)R^5$ (where R^5 is an optionally substituted alkyl, alkenyl or alkynyl group)] or a functional analogue thereof where it is
15 desired to obtain compounds wherein R^5 is a hydrogen atom, in the presence of a base, for example an alkoxide such as sodium methoxide in a solvent, for example a protic solvent, such as an alcohol, e.g. methanol at an elevated temperature, e.g. up to around the reflux temperature.

20

In a further process according to the invention, a compound of formula (1) may be prepared by displacement of a leaving atom or group in a pyrimidine of formula (6):



25

(6)


[where L is a leaving atom or group as defined above] with an aniline of formula (4).

- 30 The reaction may be performed at an elevated temperature, for example the reflux temperature, where necessary in the presence of a solvent, for example a ketone such as acetone, an alcohol such as ethanol or 2-

ethoxyethanol or an aromatic hydrocarbon such as toluene, optionally in the presence of a base, for example an organic amine such as triethylamide or pyridine, or an acid, for example an inorganic acid such as hydrochloric acid.

5

Intermediate pyrimidines of formula (6) may be prepared by displacement of a leaving group from a pyrimidine of formula (7):



10


[where L^1 is a leaving atom or group as described above for the group L] using a nucleophilic reagent R^7XH . The reaction may be performed as just described in relation to the preparation of compounds of formula (1) from the intermediate pyrimidines of formula (6).

15

The pyrimidines of formula (7) and the nucleophilic reagents R^7XH are either known compounds or may be prepared using methods analogous to those used for the preparation of the known compounds.

20

In another process according to the invention, a compound of formula (1) wherein X is a $-C(O)-$ group may be prepared by treating a carboxamide of formula (8):




[where X^3 is a carboxamide group] with a reagent $R^7M\text{Hal}^1$ (where M is a metal atom such as an zinc atom and Hal^1 is a halogen atom such as a
 5 bromine atom).

The group X^3 in intermediates of formula (8) may be for example a group $-\text{CONR}^{15}\text{R}_{16}$ or $-\text{CON}(\text{R}^{15})(\text{OR}^{16})$.

10 The reaction may be performed in a solvent such as an ether, e.g. a cyclic ether such as a tetrahydrofuran, at a low temperature, e.g. around -78°C .

Intermediate carboxamides of formula (8) may be prepared by reaction of
 15 the corresponding acid of formula (9):



or a reactive derivative thereof with a reagent $\text{R}^{15}\text{R}^{16}\text{NH}$ or $\text{R}^{15}\text{R}^{16}\text{ONH}$ in the presence of a base e.g. an organic amine such as triethylamine in a

solvent such as dichloromethane at a low temperature, e.g. around -20° to 0°C.

5 Intermediate acids of formula (9) may be prepared by heating the corresponding nitrile in the presence of a base such as sodium hydroxide in a solvent such as ethanol. The nitrile starting material may be prepared by heating 2-chloro-4-cyano-pyrimidine with the appropriate aniline in the presence of a base such as triethylamine in a solvent such as ethanol at the reflux temperature.

10

Acids of formula (9) may also be used to generate compounds of formula (1) wherein X is a -NHC(O)O- group by reaction with an azide, for example diphenylphosphorylazide, and an alcohol R⁷OH in the presence of a base such as triethylamine at an elevated temperature, e.g. the reflux temperature.

15

Compounds of formula (1) may also be prepared by interconversion of other compounds of formula (1) and it is to be understood that the invention extends to such interconversion processes. Thus, for example, standard substitution approaches employing for example alkylation, arylation, acylation, thioacylation, sulphonylation, formylation or coupling reactions may be used to add new substituents to and/or extend existing substituents in compounds of formula (1). Alternatively existing substituents in compounds of formula (1) may be modified by for example oxidation, reduction or cleavage reactions to yield other compounds of formula (1).

20

25

The following describes in general terms a number of approaches which can be employed to modify existing R¹, R², R³, R⁴, R⁵ and/or R⁶ groups in compounds of formula (1). It will be appreciated that each of these reactions may only be possible where an appropriate functional group exists in a compound of formula (1). Equally, any of the following reactions may be used to generate appropriately substituted intermediates of formulae (2), (4), (5), (6) and (7) for use in the preparation of compounds of formula (1).

30

35

Thus, for example alkylation or arylation of a compound of formula (1) may be achieved by reaction of the compound with a reagent R^4L , $AlkL$ or ArL , where R^4 , Alk , Ar and L are as previously defined.

- 5 The alkylation or arylation reaction may be carried out in the presence of a base, e.g. an inorganic base such as a carbonate, e.g. caesium or potassium carbonate, an alkoxide, e.g. potassium t-butoxide, or a hydride, e.g. sodium hydride, in a dipolar aprotic solvent such as an amide, e.g. a substituted amide such as dimethylformamide or an ether, e.g. a cyclic
10 ether such as tetrahydrofuran, at around $0^{\circ}C$ to around $40^{\circ}C$.

In a variation of this process the leaving group L may be alternatively part of the compound of formula (1) and the reaction performed with an appropriate nucleophilic reagent in a solvent such as an alcohol, e.g.
15 ethanol, at an elevated temperature, e.g. the reflux temperature.

In another general example of an interconversion process, a compound of formula (1) may be acylated or thioacylated. The reaction may be performed for example with an acyl halide or anhydride in the presence of
20 a base, such as a tertiary amine e.g. triethylamine in a solvent such as a halogenated hydrocarbon, e.g. dichloromethane or carbon tetrachloride, or an alcohol, e.g. methanol at for example ambient temperature, or by reaction with a thioester in an inert solvent such as tetrahydrofuran at a low temperature such as around $0^{\circ}C$. The reaction is particularly suitable
25 for use with compounds of formula (1) containing primary or secondary amino groups.

In a further general example of an interconversion process, a compound of formula (1) may be formylated, for example by reaction of the compound
30 with a mixed anhydride $HCOOCOCH_3$ or with a mixture of formic acid and acetic anhydride.

Compounds of formula (1) may be prepared in another general interconversion reaction by sulphonylation, for example by reaction of the
35 compound with a reagent $AlkS(O)_2L$, or $ArS(O)_2L$ in the presence of a base, for example an inorganic base such as sodium hydride in a solvent

such as an amide, e.g. a substituted amide such as dimethylformamide at for example ambient temperature. The reaction may in particular be performed with compounds of formula (1) possessing a primary or secondary amino group.

5

In further examples of interconversion reactions according to the invention compounds of formula (1) may be prepared from other compounds of formula (1) by modification of existing functional groups in the latter.

- 10 Thus in one example, ester groups $-\text{CO}_2\text{Alk}^1$ in compounds of formula (1) may be converted to the corresponding acid $[-\text{CO}_2\text{H}]$ by acid- or base-catalysed hydrolysis or by catalytic hydrogenation depending on the nature of the group Alk^1 . Acid- or base-catalysed hydrolysis may be achieved for example by treatment with an organic or inorganic acid, e.g.
- 15 trifluoroacetic acid in an aqueous solvent or a mineral acid such as hydrochloric acid in a solvent such as dioxan or an alkali metal hydroxide, e.g. lithium hydroxide in an aqueous alcohol, e.g. aqueous methanol. Catalytic hydrogenation may be carried out using for example hydrogen in the presence of a metal catalyst, for example palladium on a support such
- 20 as carbon in a solvent such as an ether, e.g. tetrahydrofuran or an alcohol, e.g. methanol. Similarly, base-catalysed hydrolysis with for example an alkali metal hydroxide such as sodium hydroxide in a solvent such as an alcohol e.g. ethanol may be used to convert a $>\text{NSO}_2\text{Alk}$ or $>\text{NSO}_2\text{Ar}$ group to a $>\text{N-H}$ group.

25

In a second example, $-\text{OAlk}^2$ [where Alk^2 represents an alkyl group such as a methyl group] groups in compounds of formula (1) may be cleaved to the corresponding alcohol $[-\text{OH}]$ by reaction with boron tribromide in a solvent such as a halogenated hydrocarbon, e.g. dichloromethane at a low

30 temperature, e.g. around -78°C .

- Alcohol $[-\text{OH}]$ groups may also be obtained by hydrogenation of the corresponding $-\text{OCH}_2\text{Ar}$ group using for example hydrogen in the presence of a metal catalyst, for example palladium on a support such as
- 35 carbon in a solvent such as ethanol in the presence of ammonium formate. In another example, $-\text{OH}$ groups may be generated from the

corresponding ester [-CO₂Alk] by reduction using for example a complex metal hydride such as lithium aluminium hydride.

5 In a further example, alcohol -OH groups in compounds of formula (1) may be converted to a corresponding -OAlk or -OAr group by coupling with a reagent AlkOH or ArOH in a solvent such as tetrahydrofuran in the presence of a phosphine, e.g. triphenylphosphine and an activator such as diethyl-, diisopropyl-, or dimethylazodicarboxylate.

10 In another example of an interconversion reaction, amines of formula (1) may be alkylated using a reductive alkylation process employing an aldehyde and a borohydride, for example sodium triacetoxyborohydride, in a solvent such as dichloromethane, in the presence of an acid such as acetic acid at around ambient temperature.

15 Aminosulphonylamino [-NH₂SO₂NH₂] groups in compounds of formula (1) may be obtained, in another example, by reaction of a corresponding amine [-NH₂] with sulphamide in the presence of an organic base such as pyridine at an elevated temperature, e.g. the reflux temperature.

20 In a further example, amine [-NH₂] groups in compounds of formula (1) may be obtained by hydrolysis from a corresponding imide by reaction with hydrazine in a solvent such as an alcohol, e.g. ethanol at ambient temperature.

25 In another example, a nitro [-NO₂] group may be reduced to an amine [-NH₂], for example by catalytic hydrogenation as just described, or by chemical reduction using for example a metal, e.g. tin or iron, in the presence of an acid such as hydrochloric acid, optionally in a solvent such as an alcohol, e.g. methanol.

30 In a further example of an interconversion process, a tetrazole substituent may be obtained from the corresponding nitrile by treatment of the latter with an azide, e.g. sodium azide, in a solvent such as a substituted amine, e.g. dimethylformamide at an elevated temperature.

N-oxides of compounds of formula (1) may be prepared for example by oxidation of the corresponding nitrogen base using an oxidising agent such as hydrogen peroxide in the presence of an acid such as acetic acid, at an elevated temperature, for example around 70°C to 80°C, or
5 alternatively by reaction with a peracid such as peracetic acid or 3-chloroperoxybenzoic acid in a solvent, e.g. dichloromethane, at ambient temperature.

Where salts of compounds of formula (1) are desired, these may be prepared by conventional means, for example by reaction of a compound
10 of formula (1) with an appropriate acid or base in a suitable solvent or mixture of solvents, e.g. an organic solvent such as an ether, e.g. diethylether, or an alcohol, e.g. ethanol.

15 The following Examples illustrated the invention. In the Examples all ¹Hnmr were run at 300MHz unless specified otherwise. All temperatures are in °C. The following abbreviations are used: DMSO - dimethylsulphoxide; DMF - dimethylformamide; THF - tetrahydrofuran.

20 **EXAMPLE 1**

4-Phenyl-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine

To a solution of 3,4,5-(trimethoxyphenyl)guanidine nitrate (1.89g, 6.57mmol) and 3-dimethylamino-1-phenyl-2-propen-1-one (1.15g, 6.57mmol) in propan-2-ol (20ml) was added powdered sodium hydroxide
25 (289mg, 7.23mmol) and the mixture refluxed for 18h. On cooling (0°) the resulting precipitate was collected and washed with propan-2-ol and water, and then subjected to column chromatography [silica 1% methanol-CH₂CH₂] to afford the title compound (730mg) after recrystallisation from ethyl acetate as a yellow solid m.p. 146°. δ_H (CDCl₃) 3.85 (3H, s), 3.91
30 (6H, s), 7.07 (2H, s), 7.17 (1H, d, J 5.2Hz), 7.18 (1H, s, NH), 7.46-7.50 (3H, m), 8.10 (2H, m), and 8.46 (1H, d, J 5.2Hz).

The guanidine starting material was prepared by heating a mixture of 3,4,5-trimethoxyaniline (5.49g, 30.0mmol), cyanamide [Aldrich, 50%
35 solution in water w/v] (3.50ml, 45.0mmol) and concentrated nitric acid (2.10ml, 300mmol) in ethanol (30ml). The solid which formed on cooling

to room temperature was collected by filtration, washed with ethanol and dried *in vacuo* to give the desired product (4.60g) as a grey solid m.p. 187°. δ_H (d^6 DMSO) 3.65 (3H, s), 3.77 (6H, s), 6.54 (2H, s), 7.27 (4H, br s), and 9.46 (1H, s).

5

The following compounds of Examples 2-35 were prepared in a similar manner using the above guanidine starting material and the appropriate enaminone:

10 **EXAMPLE 2****4-(3-Pyridyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

from 3,4,5-(trimethoxyphenyl)guanidine nitrate (1.33g, 4.60mmol), 3-dimethylamino-1-(3-pyridyl)-2-propen-1-one (812mg, 4.60mmol) and sodium hydroxide (203mg, 5.07mmol) to give the title compound (290mg) as a green solid m.p. 155°. δ_H ($CDCl_3$) 3.63 (3H, s), 3.78 (6H, s), 7.28 (2H, s), 7.47 (1H, d, \downarrow 5.1Hz), 7.56 (1H, m), 8.49 (1H, m), 8.58 (1H, d, \downarrow 5.1Hz), 8.71 (1H, dd, \downarrow 4.8, 1.5Hz), 9.35 (1H, d, \downarrow 2.0Hz) and 9.61 (1H, s).

20 **EXAMPLE 3****4-(4-Pyridyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

from 3,4,5-(trimethoxyphenyl)guanidine nitrate (1.44g, 5.0mmol), 3-dimethylamino-1-(4-pyridyl)-2-propene-1-one (880mg, 5.0mmol) and sodium hydroxide (220mg, 5.5mmol) to give the title compound (765mg) as a green solid m.p. 205°. δ_H (d^6 DMSO) 3.63 (3H, s), 3.79 (6H, s), 7.27 (2H, s), 7.49 (1H, d, \downarrow 5.1Hz), 8.08 (2H, dd, \downarrow 4.5, 1.6Hz), 8.64 (1H, d, \downarrow 5.1Hz), 8.76 (2H, dd, \downarrow 4.5, 1.6Hz), and 9.66 (1H, br s).

30 **EXAMPLE 4****4-(2-Furyl)-N-3,4,5-(trimethoxyphenyl)-2-pyrimidineamine**

from 3,4,5-(trimethoxyphenyl)guanidine nitrate (1.66g, 5.75mmol), 3-dimethylamino-1-(2-furyl)-2-propen-1-one (950mg, 5.75mmol) and sodium hydroxide (253mg, 6.33 mmol) to give the title compound (350mg) as a yellow solid m.p. 139°. δ_H ($CDCl_3$) 3.83 (3H, s), 3.90 (6H, s), 6.56 (1H, m), 7.01 (2H, s), 7.06 (1H, d, \downarrow 5.1Hz), 7.18 (1H, m), 7.25 (1H, s), 7.58 (1H, d, \downarrow 2.3 Hz), and 8.42 (1H, d, \downarrow 5.1Hz).

35

EXAMPLE 5**4-(3,4,5-Trimethoxyphenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidine-amine**

from 3,4,5-(trimethoxyphenyl)guanidine nitrate (1.08g, 3.77mmol), 3-dimethylamino-1-(3,4,5-trimethoxyphenyl)-2-propen-1-one (1.0g, 3.77mmol) and sodium hydroxide (166mg) to give the title compound (67mg) as a colourless solid m.p. 187°. δ_H (CDCl₃) 3.83 (3H, s), 3.88 (6H, s), 3.92 (3H, s), 3.94 (6H, s), 6.99 (2H, s), 7.09 (1H, d, \downarrow 5.2Hz), 7.23 (1H, br s), 7.32 (2H, s) and 8.45 (1H, d, \downarrow 5.2Hz).

EXAMPLE 6**4-(5-(2,3-Dihydrobenzofuryl))-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

from 3,4,5-(trimethoxyphenyl)guanidine nitrate (1.19g, 4.14mmol), 1-(5-(2,3-dihydrobenzofuryl)-3-dimethylamino-2-propen-1-one (900mg, 4.14mmol) and sodium hydroxide (182mg, 4.56mmol) to give the title compound (450mg) as a yellow solid m.p. 160°. δ_H (CDCl₃) 3.26 (2H, t, \downarrow 8.7Hz), 3.83 (3H, s), 3.90 (6H, s), 4.66 (2H, t, \downarrow 8.7Hz), 6.86 (1H, d, \downarrow 8.3Hz), 7.06 (3H, m), 7.17 (1H, s), 7.88 (1H, d, \downarrow 8.3Hz), 8.02 (1H, s), and 8.38 (1H, d, \downarrow 5.3Hz).

EXAMPLE 7**4-(1-Phenylsulphonylindol-3-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

from 3,4,5-(trimethoxyphenyl)guanidine nitrate (820mg, 2.84mmol), 3-dimethylamino-1-(1-phenylsulphonylindol-3-yl)-2-propen-1-one (1.0g, 2.84mmol), and sodium hydroxide (125mg, 3.12mmol) to give the title compound (380mg) as a yellow solid m.p. 202°. δ_H (d⁶DMSO) 3.30 (3H, s), 3.63 (6H, s), 7.19 (2H, s), 7.33 (1H, t, \downarrow 7.3Hz), 7.44 (2H, m), 7.61 (2H, m), 7.71 (1H, m), 7.99 (1H, d, \downarrow 8.3Hz), 8.09 (2H, d, \downarrow 7.4Hz), 8.49 (1H, d, \downarrow 5.2Hz), 8.60 (1H, d, \downarrow 7.9Hz), 8.74 (1H, s) and 9.46 (1H, s).

EXAMPLE 8**4-(2-Thiazolyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

from 3,4,5-(trimethoxyphenyl)guanidine nitrate (1.27g, 4.40mmol), 3-dimethylamino-1-(2-thiazolyl)-2-propen-1-one (802mg, 4.40mmol), and

sodium hydroxide (194mg, 4.84mmol) to give the title compound (520mg) as a green solid m.p. 159°. δ_H (d^6 DMSO) 3.63 (3H, s), 3.82 (6H, s), 7.24 (2H, s), 7.45 (1H, d, J 4.9Hz), 8.07 (1H, d, J 3.0Hz), 8.62 (1H, d, J 4.8Hz) and 9.70 (1H, s).

5

EXAMPLE 9**4-(3-Thienyl)-N-(3,4,5-Trimethoxyphenyl)-2-pyrimidineamine**

from 3,4,5-(trimethoxyphenyl)guanidine nitrate (1.27g, 4.41mmol), 3-dimethylamino-1-(3-thienyl)-2-propen-1-one (0.80g, 4.41mmol) to give the title compound (365mg) as a yellow solid m.p. 163°. δ_H (d^6 DMSO) 3.62 (3H, s), 3.78 (6H, s), 7.26-7.27 (3H, m), 7.68-7.71 (1H, m), 7.78 (1H, d, J 4.9Hz), 8.34-8.36 (1H, m), 8.47 (1H, d, J 5.0Hz) and 9.44 (1H, s).

10

EXAMPLE 10**4-(2-Naphthyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

from 3,4,5-(trimethoxyphenyl)guanidine nitrate (1.02g, 3.55mmol), 3-dimethylamino-1-(2-naphthyl)-2-propen-1-one (800mg, 3.55mmol) and sodium hydroxide (156mg, 3.90mmol) to give the title compound (310mg) as a yellow solid m.p. 156°. δ_H (d^6 DMSO) 3.64 (3H, s), 3.82 (6H, s), 7.36 (2H, s), 7.53 (1H, d, J 6.2Hz), 7.58-7.61 (2H, m), 7.97-8.07 (3H, m), 8.29 (1H, d, J 8.5Hz), 8.58 (1H, d, J 5.1Hz), 8.76 (1H, s), and 9.58 (1H, br s).

15

20

EXAMPLE 11**4-(3-Nitrophenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

from 3,4,5-(trimethoxyphenyl)guanidine nitrate (3.93g, 13.63mmol), 3-dimethylamino-1-(3-nitrophenyl)-2-propen-1-one (3.0g, 13.63mmol) and sodium hydroxide (545mg, 13.63mmol) to give the title compound (250mg) as a yellow solid. m.p. 184-185°. MS m/z 383 (M+H)⁺.

30

EXAMPLE 12**4-(4-Nitrophenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

from 3,4,5-(trimethoxyphenyl)guanidine nitrate (1.3g, 4.5mmol), 3-dimethylamino-1-(4-nitrophenyl)-2-propen-1-one (1.0g, 4.5mmol) and sodium hydroxide (200mg) to give the title compound (550mg) as an orange solid m.p. 196°. MS m/z 383 (M+H)⁺.

35

EXAMPLE 13**4-(3-Bromophenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

from 3,4,5-(trimethoxyphenyl)guanidine nitrate (1.63g, 5.66mmol), 1-(3-bromophenyl)-3-dimethylamino-2-propen-1-one (1.5g, 5.66mmol) and sodium hydroxide (250mg, 6.23mmol) to give the title compound (785mg) as a yellow solid m.p. 145°. MS m/z 418 (M+H)⁺.

EXAMPLE 14**4-(Pyrrol-2-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

from 3,4,5-(trimethoxyphenyl)guanidine nitrate (1.15g, 4.0mmol), 3-dimethylamino-1-(pyrrol-2-yl)-2-propen-1-one (656g, 4.0mmol) and sodium hydroxide (176mg, 4.4mmol) to give the title compound (10mg) as a yellow solid m.p. 150°. MS m/z 327 (M+H)⁺.

EXAMPLE 15**4-(3,4-Methylenedioxyphenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

from 3,4,5-(trimethoxyphenyl)guanidine nitrate (1.15g, 4.0mmol), 3-dimethylamino-1-(3,4-methylenedioxyphenyl)-2-propen-1-one (8.76g, 4.0mmol) and sodium hydroxide (176mg) to give the title compound (160mg) as a yellow solid m.p. 180°. MS m/z 382 (M+H)⁺.

EXAMPLE 16**4-(Pyrazin-2-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

from 3,4,5-(trimethoxyphenyl)guanidine nitrate (1.15g, 4.0mmol), 3-dimethylamino-1-(pyrazin-2-yl)-2-propen-1-one (708mg, 4.0mmol) and sodium hydroxide (176mg) to give the title compound (254mg) as a yellow solid. m.p. 181-182°. MS m/z 340 (M+H)⁺.

EXAMPLE 17**4-(tert-Butyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

from 3,4,5-(trimethoxyphenyl)guanidine nitrate (1.2g, 4.2mmol), 1-tert-butyl-3-dimethylamino-2-propen-1-one (650mg, 4.2mmol) and sodium hydroxide (185mg) to give the title compound (90mg) as a white solid m.p. 122°. MS m/z 318 (M+H)⁺.

EXAMPLE 18**4-(1,2,3,4-Tetrahydronaphthalen-6-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

5 from 3,4,5-(trimethoxyphenyl)guanidine nitrate (1.0g, 4.35mmol), 3-dimethylamino-1-(1,2,3,4-tetrahydronaphthalen-6-yl)-2-propen-1-one (801mg, 3.5mmol) and sodium hydroxide (155mg, 3.85mmol) to give the title compound (120mg) as a yellow solid m.p. 150-151°. MS m/z 392 (M+H)⁺.

10 EXAMPLE 19**4-(2,2-Dimethyl-3,4-dihydro-2H-benzo[b]oxin-6-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

15 from 3,4,5-(trimethoxyphenyl)guanidine nitrate (1.15g, 4.0mmol), 3-dimethylamino-1-(2,2-dimethyl-3,4-dihydro-2H-benzo[b]oxin-6-yl)-2-propen-1-one (1.04g, 4.0mmol) and sodium hydroxide (180mg, 4.4mmol) to give the title compound (120mg) as a pale yellow solid m.p. 149-150°. MS m/z 422.3 (M+H)⁺.

EXAMPLE 20**20 4-(Benzothiazol-2-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

from 3,4,5-(trimethoxyphenyl)guanidine nitrate (1.72g, 6.0mmol), 1-(benzothiazol-2-yl)-3-dimethylamino-2-propen-1-one (1.39g, 6.0mmol) and sodium hydroxide (270mg, 6.6mmol) to give the title compound (420mg) as a yellow solid m.p. 182-183°. MS m/z 395 (M+H)⁺.

25

EXAMPLE 21**4-(2-Nitrothien-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

30 from 3,4,5-(trimethoxyphenyl)guanidine nitrate (1.27g, 4.4mmol), 3-dimethylamino-1-(2-nitrothien-5-yl)-2-propen-1-one (1.0g, 4.4mmol) and sodium hydroxide (186mg, 4.9mmol) to give the title compound (540mg) as a dark red solid m.p. 182-183°. MS m/z 389 (M+H)⁺.

EXAMPLE 22**4-(3-Methoxyphenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

35 from 3,4,5-(trimethoxyphenyl)guanidine nitrate (1.41g, 4.88mmol), 3-dimethylamino-1-(3-methoxyphenyl)-2-propen-1-one (1.0g, 4.88mmol) and

sodium hydroxide (216mg, 5.4mmol) to give the title compound (275mg) as a yellow solid m.p. 155-156°. MS m/z 368 (M+H)⁺.

EXAMPLE 23

- 5 **4-(2-Pyridyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**
from 3,4,5-(trimethoxyphenyl)guanidine nitrate (1.99g, 6.91mmol), 3-dimethylamino-1-(2-pyridyl)-2-propen-1-one (1.22g, 6.91mmol) and sodium hydroxide (276mg, 6.9mmol) to give the title compound (158mg) as a yellow solid m.p. 185°. MS m/z 339 (M+H)⁺.

10

EXAMPLE 24

- 4-(4-*tert*-Butoxycarbonylaminophenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**
from 3,4,5-(trimethoxyphenyl)guanidine nitrate (10.9g, 37.8mmol), 1-(4-*tert*-butoxycarbonylaminophenyl)-3-dimethylamino-2-propen-1-one (10.0g, 34.4mmol) and sodium hydroxide (1.52g, 37.8mmol) to give the title compound (5.4g) as a yellow solid m.p. 158-159°. MS m/z 453 (M+H)⁺.

15

EXAMPLE 25

- 4-(3-Hydroxyphenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**
from 3,4,5-(trimethoxyphenyl)guanidine nitrate (4.72g, 16.4mmol), 1-(3-*tert*-butyldimethylsilyloxyphenyl)-3-dimethylamino-2-propen-1-one (5.0g, 16.4mmol) and sodium hydroxide (655mg, 16.4mmol) to give the title compound (2.0g) as a light yellow solid m.p. 191-192°. MS m/z 354 (M+H)⁺.

25

EXAMPLE 26

- 4-(5-Thiazolyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**
from 3,4,5-(trimethoxyphenyl)guanidine nitrate (634mg, 2.2mmol), 3-dimethylamino-1-(5-thiazolyl)-2-propen-1-one (330g, 1.81mmol) and sodium hydroxide (88mg, 2.2mmol) to give the title compound (42mg) as a yellow solid m.p. 171°. MS m/z 345 (M+H)⁺.

30

35

EXAMPLE 27**4-(4-Cyanophenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

from 3,4,5-(trimethoxyphenyl)guanidine nitrate (17.0g, 59.0mmol), 1-(4-cyanophenyl)-3-dimethylamino-2-propen-1-one (11.82g, 58.0mmol) and
5 sodium hydroxide (2.36mg, 59.0mmol) to give the title compound (11.66g) as a green solid m.p. 183°. MS m/z 363 (M+H)+.

EXAMPLE 28**4-(4-Ethoxycarbonylphenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

10 from 3,4,5-(trimethoxyphenyl)guanidine nitrate (11.66g, 40.5mmol), 3-dimethylamino-1-(4-ethoxycarbonylphenyl)-2-propen-1-one (10.0g, 40.5mmol) and sodium hydroxide (1.62g, 40.5mmol) to give the title compound (3.21g) as a yellow solid m.p. 181-182°. MS m/z 410 (M+H)+.

15

EXAMPLE 29**4-(1-Naphthyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

from 3,4,5-(trimethoxyphenyl)guanidine nitrate (2.94g, 10.22mmol), 3-dimethylamino-1-(1-naphthyl)-2-propen-1-one (2.30g, 10.22mmol) and
20 sodium hydroxide (409mg, 10.22mmol) to give the title compound (750mg) as a yellow solid m.p. 130-133°. MS m/z 444 (M+H)+.

EXAMPLE 30**4(4,5-Dimethylthiazol-2-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

25 from 3,4,5-(trimethoxyphenyl)guanidine nitrate (4.99g, 6.91mmol), 3-dimethylamino-1-(4,5-dimethylthiazol-2-yl)-2-propen-1-one (3.64g, 17.3mmol) and sodium hydroxide (720mg, 18mmol) to give the title compound (490mg) as a yellow solid m.p. 207°. MS m/z 373 (M+H)+.

30

EXAMPLE 31**4-(4-Methoxyphenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

from 3,4,5-(trimethoxyphenyl)guanidine nitrate (8.16g, 28.3mmol), 3-dimethylamino-1-(4-methoxyphenyl)-2-propen-1-one (5.5g, 28.3mmol) and
35 sodium hydroxide (1.13g, 28.3mmol) to give the title compound (3.4g) as an orange solid m.p. 148-150°. MS m/z 368 (M+H)+.

EXAMPLE 32

4-(4-Benzyloxyphenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine
from 3,4,5-(trimethoxyphenyl)guanidine nitrate (9.0g, 31.25mmol), 1-(4-benzyloxyphenyl)-3-dimethylamino-2-propen-1-one (8.88g, 31.25mmol)
5 and sodium hydroxide (1.25g, 31.25mmol) to give the title compound (11.0g) as a yellow solid m.p. 171-172°. MS m/z 444(M+H)⁺.

EXAMPLE 33

4-(5-(2-Hydroxyethyl)-4-methylthiazol-2-yl)-N-(3,4,5-trimethoxy-phenyl)-2-pyrimidineamine

10 from 3,4,5-(trimethoxyphenyl)guanidine nitrate (1.22g, 4.23mmol), 1-(5-(2-*tert*-butyldimethylsilyloxyethyl)-4-methylthiazol-2-yl)-3-dimethylamino-2-propen-1-one (1.50g, 4.23mmol) and sodium hydroxide (200mg, 5.0mmol) to give the title compound (210mg) as an orange solid m.p. >190°
15 (decomp). MS m/z 403 (M+H)⁺.

EXAMPLE 34

4-(4-Bromophenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine

20 from 3,4,5-(trimethoxyphenyl)guanidine nitrate (2.26g, 7.80mmol), 1-(4-bromophenyl)-3-dimethylamino-2-propen-1-one (2.0g, 7.80mmol) and sodium hydroxide (346mg, 8.60mmol) to give the title compound (1.0g) as a green solid m.p. 179°. MS m/z 416 (M+H)⁺.

EXAMPLE 35

25 **4-(2-Chloropyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidine-amine**

30 from 3,4,5-(trimethoxyphenyl)guanidine nitrate (16.42g, 22.3mmol), 1-(2-chloropyridin-5-yl)-3-dimethylamino-2-propen-1-one (4.70g, 22.3mmol) and sodium hydroxide (895mg) to give the title compound (1.43g) as a yellow solid m.p. 191-192°. MS m/z 373.2 (M+H)⁺.

EXAMPLE 36

4-(1-H-Indol-3-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine

35 The compound of Example 7 (400mg, 0.78mmol) was suspended in ethanol (5ml) and 4M sodium hydroxide (5ml) and heated at reflux for 4h. On cooling a precipitate formed which was collected by filtration and

washed with water to give the title compound (79mg) as a yellow solid m.p. 197°. MS m/z 377 (M+H)⁺.

EXAMPLE 37

5 **N-Methyl-4-(4-pyridyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

A solution of the compound of Example 3 (650mg, 1.92mmol) in DMF (5ml) was added dropwise to a stirred suspension of 60% NaH (76mg, 1.92mmol) in DMF (5ml), cooled to 0°C. After 0.5h iodomethane (0.13ml, 2.11mmol) was added dropwise and the resulting mixture was left to warm
10 to room temperature over 20h. The reaction was concentrated under reduced pressure and the resulting residue subjected to column chromatography to give the title compound (159mg) as a pale green solid m.p. 139°. MS m/z 353 (M+H)⁺.

15 **EXAMPLE 38**

4-(2-Aminothien-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine

A solution of the compound of Example 21 (380mg, 0.98mmol) was treated with ammonium formate (200mg, 6.34mmol) and 10% palladium on carbon (200mg) at 65° for 24h. On cooling, the catalyst was filtered off
20 on a pad of celite and the filtrate concentrated under reduced pressure to give the title compound (100mg) as a green solid m.p. 161-162°. MS m/z 389 (M+H)⁺.

The compounds of Examples 39 and 40 were prepared in a similar
25 manner to the compound of Example 38:

EXAMPLE 39

4-(4-Aminophenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine

from the compound of Example 12 (800mg, 2.09mmol), ammonium formate (660mg, 10.5mmol) and 10% palladium on carbon (80mg) to give
30 the title compound (253mg) as a yellow solid m.p. 195°. MS m/z 353 (M+H)⁺.

EXAMPLE 40**4-(3-Aminophenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

from the compound of Example 11 (150mg, 0.39mmol), ammonium formate (150mg, 2.38mmol) and 10% palladium on carbon (100mg) to give the title compound (120mg) as a yellow solid m.p. 166-167°. MS m/z 353 (M+H)⁺.

EXAMPLE 41**4-(3-Acetamidophenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

To a solution of the compound of Example 40 (0.25g, 0.71mmol) in dry CH₂Cl₂ (40ml) and triethylamine (0.11ml) was added acetyl chloride (0.06ml, 0.78mmol) and the resulting mixture stirred at room temperature for 4h. After washing with water (100ml) and saturated aqueous NaHCO₃ (100ml), the organic layer was dried (MgSO₄) and concentrated under reduced pressure. The residue was subjected to column chromatography [silica-ethyl acetate] to give after recrystallisation from ethyl acetate-hexane the title compound (163mg) as a yellow solid. m.p. 162-164°. MS m/z 395(M+H)⁺.

The compounds of Examples 42-44 were prepared in a similar manner to the compound of Example 41:

EXAMPLE 42**4-(4-Acetamidophenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

from the compound of Example 39 (300mg, 0.85mmol), triethylamine (0.13ml, 0.94mmol) and acetyl chloride (0.07ml, 0.94mmol) to give the title compound (178mg) as a yellow solid m.p. 239°. MS m/z 395 (M+H)⁺.

EXAMPLE 43**N-(3-(2-(3,4,5-Trimethoxyphenylamino)pyrimidin-4-yl)phenyl)-5-methylisoxazole-3-carboxamide**

from the compound of Example 40 (150mg, 0.43mmol), 5-methylisoxazole-3-carbonyl chloride (68mg, 0.47mmol) and triethylamine (0.1ml, 0.7mmol) to give the title compound (7mg) as a yellow solid m.p. 173-175°. MS m/z 462 (M+H)⁺.

EXAMPLE 44**N-(3-(2-(3,4,5-Trimethoxyphenylamino)pyrimidin-4-yl)phenyl)pyridine-3-carboxamide**

from the compound of Example 40 (250mg, 0.71mmol), nicotinoyl chloride hydrochloride (139mg, 0.78mmol), and triethylamine (0.2ml, 2.3mmol) to give the title compound (155mg) as a yellow solid m.p. 140-143°. MS m/z 458 (M+H)⁺.

EXAMPLE 45

10 **4-(3-(3-Phthalimidopropoxy)phenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

To a suspension of caesium carbonate (462mg, 1.4mmol) in dry DMF (10ml) under a nitrogen atmosphere were added the compound of Example 25 (500mg, 1.4mmol) and 3-bromopropylphthalimide (380mg, 1.4mmol). The resulting mixture was stirred at room temperature for 4.5h, followed by the addition of water (50ml). The resulting precipitate was collected, washed with diethyl ether and recrystallised from ethyl acetate-hexane to give the title compound (620mg) as an off-white solid m.p. 209°. MS m/z 541 (M+H)⁺.

20

The following compound was prepared in a similar manner:

EXAMPLE 46**4-(3-Propoxyphenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

from the compound of Example 25 (350mg, 0.99mmol), iodopropane (168mg, 0.99mmol) and caesium carbonate (322mg, 0.99mmol) to give the title compound (187mg) as an off-white solid m.p. 107°. MS m/z 396 (M+H)⁺.

EXAMPLE 47

30 **4-(3-(2-Hydroxyethoxy)phenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

To a suspension of 60% NaH (dispersion in oil) (62mg, 1.6mmol) in DMF (15ml) under a nitrogen atmosphere were added the compound of Example 25 (500mg, 1.4mmol) and ethylene carbonate (137mg, 1.6mmol). The resulting mixture was heated at 130° for 6h. On cooling the reaction was diluted with water (30ml), extracted with CH₂Cl₂

(3x25ml), dried (MgSO₄) and concentrated under reduced pressure. The resulting residue was subjected to column chromatography [silica-ethyl acetate] to give the title compound (260mg) as a yellow solid m.p. 70°. MS m/z 398 (M+H)⁺.

5

EXAMPLE 48**4-(3-(3-Aminopropoxy)phenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine dihydrochloride**

The compound of Example 45 (500mg, 0.9mmol) was suspended in ethanol (30ml) containing hydrazine monohydrate (0.14ml, 2.8mmol) and the resulting mixture heated at reflux for 18h. On cooling the resulting precipitate was filtered off and the filtrate concentrated under reduced pressure. The residue was suspended in CH₂Cl₂ (30ml), extracted with 2N hydrochloric acid (2x30ml). Combined acid layers were taken to pH 11 with 6N NaOH and extracted with CH₂Cl₂ (3x30ml), the organic layers were dried (MgSO₄), and concentrated under reduced pressure. The residue was dissolved in ethanol (10ml) and the solution saturated with HCl (g). The resulting precipitate was collected to give the title compound (137mg) as an orange solid m.p. 236°. δ H (d⁶-DMSO) 9.62 (1H, s), 8.54 (1H, d, \downarrow 5.2Hz), 8.05 (2H, br s), 7.75 (2H, m), 7.45 (2H, m), 7.29 (2H, s), 7.14 (1H, d, \downarrow 7.3Hz), 4.14 (2H, t, \downarrow 6.0Hz), 3.79 (6H, s), 3.62 (3H, s), 2.95 (2H, m) and 2.06 (2H, m). MS m/z 411 (M+H)⁺.

EXAMPLE 49**4-(4-Hydroxyphenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

A solution of the compound of Example 32 (1.06g, 2.37mmol) in ethanol (100ml) was treated with ammonium formate (750mg, 12.0mmol) and 10% palladium on carbon (100mg) and stirred at room temperature for 18h, followed by heating at reflux for 18h. On cooling the catalyst was removed by filtration through Celite® and the filtrate concentrated under reduced pressure. The resulting residue was recrystallised from ethyl acetate to give the title compound (760mg) as a cream solid m.p. 200-202°. MS m/z 354 (M+H)⁺.

30

EXAMPLE 50**4-(4-(3-Phthalimidopropoxy)phenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

To a suspension of 60% NaH (dispersion in oil) (55mg, 1.36mmol) in DMF (10ml) was added the compound of Example 49 (480mg, 1.36mmol) and 3-bromopropylphthalimide (365mg, 1.36mmol). The resulting mixture was stirred at room temperature for 6h. After this time the reaction was concentrated under reduced pressure, the residue dissolved in ethyl acetate (50ml), washed with water (50ml), and dried (MgSO₄). The solvent was removed under reduced pressure to give the title compound (700mg) as a buff solid. MS m/z 541 (M+H)⁺.

EXAMPLE 51**4-(4-(3-Aminopropoxy)phenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

The compound was prepared in a manner analogous to the preparation of the compound of Example 48 from the compound of Example 50 (830mg, 1.54mmol) and hydrazine monohydrate (0.23ml, 4.62mmol) to give the title compound (63mg) as a white solid m.p. 161-162°. MS m/z 411(M+H)⁺.

EXAMPLE 52**4-(4-(2-Hydroxyethoxy)phenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

The compound was prepared in a manner analogous to the preparation of the compound of Example 47 from the compound of Example 49 (250mg, 0.71mmol), ethylene carbonate (70mg, 6.78mmol) and 60% sodium hydride (dispersion in oil) (30mg, 0.78mmol) to give the title compound (140mg) as a yellow solid m.p 145-146°. MS m/z 398 (M+H)⁺.

EXAMPLE 53**4-(4-(2-N,N-Dimethylaminoethoxy)phenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

The compound of Example 49 (250mg, 0.71mmol), N,N-dimethylaminoethanol (69mg, 0.78mmol) and triphenylphosphine (205mg, 0.78mmol) were dissolved in dry THF (20ml) under a nitrogen atmosphere and the mixture stirred for 0.5h. Diethylazodicarboxylate (136mg, 0.78mmol) in

THF (10ml) was added dropwise and the reaction stirred for 48h, after which time the solvent was removed under reduced pressure. The residue was dissolved in ethyl acetate (40ml), washed with water (2x100ml), dried (MgSO₄) and the solvent evaporated. The resulting solid
5 was subjected to column chromatography [silica-ethyl acetate] to give the title compound (75mg) as white crystals m.p. 140-147°. MS m/z 425 (M+H)⁺.

The following compound was prepared in a similar manner:

10 **EXAMPLE 54**

4-(4-(3-N,N-Dimethylaminopropoxy)phenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine

from the compound of Example 49 (250mg, 0.71mmol), N,N-dimethylaminopropanol (88mg, 0.85mmol), triphenylphosphine (223mg,
15 0.85mmol) and diethylazodicarboxylate (148mg, 0.85mmol) to give the title compound (20mg) as a white solid m.p. 161-164°. MS m/z 439 (M+H)⁺.

EXAMPLE 55

20 **4-(3-(3-Hydroxypropylamino)phenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

The compound of Example 40 (250mg, 0.71mmol) and 3-bromopropanol were heated at 85° in DMF for 72h. The solvent was removed under reduced pressure and the residue subjected to column chromatography [silica-ethyl acetate] to give the title compound (33mg) as a yellow solid
25 m.p. 92-95°. MS m/z 411 (M+H)⁺.

The following compound was prepared in a similar manner:

EXAMPLE 56

30 **4-(4-(3-Hydroxypropylamino)phenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

from the compound of Example 39 (750mg, 2.13mmol) and 3-bromopropanol (3.55mg, 2.56mmol) to give the title compound (24mg) as a yellow solid m.p. 86-90°. MS m/z 411 (M+H)⁺.

EXAMPLE 57**4-(3-(3-Pyridylmethyl)aminophenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

To a solution of the compound of Example 40 (250mg, 0.71mmol) in
5 CH₂Cl₂ (20ml) was added 3-pyridinecarboxaldehyde (69mg, 0.65mmol),
sodium triacetoxyborohydride (226mg, 1.07mmol) and acetic acid (0.1ml),
and the resulting mixture stirred at ambient temperature for 48h. The
reaction was washed with water (2x100ml), dried (MgSO₄), concentrated
under reduced pressure, and the residue columned [silica-ethyl acetate] to
10 give the title compound (100mg) as a yellow solid m.p. 78-80°. MS m/z
444 (M+H)⁺.

The following compound was prepared in a similar manner:

EXAMPLE 58

15 **4-(4-(3-Pyridylmethyl)aminophenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

from the compound of Example 39 (250mg, 0.71mmol), 3-pyridine-
carboxaldehyde (69mg, 0.65mmol), sodium triacetoxyborohydride (226mg,
1.67mmol) and acetic acid (0.1ml) to give the title compound (134mg) as a
20 yellow solid m.p. 189-190°. MS m/z 444 (M+H)⁺.

EXAMPLE 59**4-(4-(Aminomethyl)phenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

25 To a suspension of LiAlH₄ (628mg, 16.56mmol) in dry THF (100ml) at 0°
was added dropwise a solution of the compound of Example 27 in THF
(10ml) and when addition was complete the reaction was heated at reflux
for 3h. On cooling a saturated solution of aqueous ammonium chloride
(25ml) was added and the organic solvent removed under reduced
30 pressure. The aqueous solution was extracted with ethyl acetate
(2x200ml) and the combined organic layers dried (MgSO₄) followed by
evaporation under reduced pressure. The residue was subjected to
column chromatography [silica-ethyl acetate] to give the title compound
(330mg) as a yellow solid m.p. 143-144°. MS m/z 367 (M+H)⁺.

EXAMPLE 60**4-(4-(1H-1,2,3,4-Tetrazol-5-yl)phenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

- 5 To a solution of the compound of Example 27 (500mg, 1.38mmol) in DMF (15ml) was added sodium azide (897mg, 13.8mmol) and ammonium chloride (738mg, 13.8mmol) and the mixture heated at 130° for 18h. On cooling, water (45ml) was added, the resulting precipitate collected and recrystallised from ethyl acetate to give the title compound (333mg) as a
- 10 yellow solid m.p. 246-247°. δ H 9.60 (1H, br s), 8.59 (1H, d, \downarrow 5.0Hz), 8.39 (2H, d, \downarrow 8.4), 8.19 (2H, d, \downarrow 8.4Hz), . 7.47 (1H, d, \downarrow 5.3Hz), 7.31 (2H, s), 3.80 (6H, s) and 3.62 (3H, s). MS m/z 406.2 (M+H)⁺.

EXAMPLE 61**4-(1-Oxopyrid-4-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

- 15 In a manner analogous to the preparation of the compound of Example 1, from 3,4,5-trimethoxyphenylguanidine nitrate (2.25g, 7.81mmol), 3-dimethylamino-1-(1-oxo-pyrid-4-yl)-2-propen-1-one (880mg, 7.81mmol) and sodium hydroxide (344mg, 8.6mmol) to give the title compound (1.2g) as a green solid m.p. 220°. MS m/z 355 (M+H)⁺.
- 20

- The 3-dimethylamino-1-(1-oxopyrid-4-yl)-2-propen-1-one used as starting material was prepared by heating a solution of 4-acetylpyridine-N-oxide (2.5g, 18.2mmol) in dimethylformamide diethylacetal (30ml) at reflux for 0.5h. On cooling the resulting solid was collected and washed with diethyl
- 25 ether to give the desired product (3.18g) as an orange solid m.p. 181°.

- The 4-acetylpyridine-N-oxide was prepared by treating a solution of 4-acetylpyridine (3.0g, 24.8mmol) in CH₂Cl₂ with 3-chloro-peroxybenzoic acid [Aldrich 57-86%] (8.4g) at room temperature for 12 h. The reaction was filtered, the filtrate concentrated under reduced pressure and the
- 30 resulting residue subjected to column chromatography [silica 10% methanol-ethyl acetate] to give the desired product (3.2g) as a white solid m.p. 101°.

EXAMPLE 62**4-(4-(Hydroxymethyl)phenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

To a suspension of LiAlH₄ (560mg, 14.64mmol) in dry THF (50ml) at 0°
5 was added dropwise to a solution of the compound of Example 28 in THF
(100ml) and the reaction heated at reflux for 18h. On cooling an aqueous
saturated solution of ammonium chloride (70ml) was added and the
organic solvent removed under reduced pressure. The aqueous solution
10 was extracted with CH₂Cl₂ (300ml), and this was dried (MgSO₄) and
evaporated under reduced pressure. The residue was subjected to
column chromatography [silica-ethyl acetate] to give title compound
(212mg) as a pale yellow solid m.p. 171-172°. MS m/z 368 (M+H)⁺.

EXAMPLE 63

15 **4-(2-(3-Hydroxypropylamino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

The compound of Example 35 (200mg, 0.54mmol) and 3-amino-1-
propanol (1.0ml, 13.05mmol) were heated at 100° for 3.5h. On cooling the
reaction was concentrated under reduced pressure, water (15ml) added
20 and the resulting precipitate collected. After washing with water, drying
under vacuum and recrystallisation from ethanol the title compound
(105mg) was obtained as a yellow solid m.p. 168-160°. δ H(d⁶ DMSO)
9.40 (1H, s), 8.83 (1H, s), 8.38 (1H, d, \downarrow 5.3Hz), 8.12 (1H, br dm, \downarrow 7.0Hz),
7.26 (2H, s), 7.22 (1H, d, \downarrow 5.2Hz), 7.16 (1H, br t, \downarrow 3.0Hz), 6.54 (1H, d, \downarrow
25 8.8Hz), 4.53 (1H, t, \downarrow 5.0Hz), 3.76 (6H, s), 3.60 (3H, s), 3.47 (2H, q, \downarrow
5.4Hz), 3.32 (2H, m) and 1.68 (2H, m). MS m/z 412 (M+H)⁺.

The compounds of Examples 64-87 were prepared in a similar manner to
the compound of Example 63 using the compound of Example 35 and the
30 amine shown:

EXAMPLE 64**4-(2-(2-Aminoethylamino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

from the compound of Example 35 (200mg, 0.54mmol) and ethylene-
35 diamine (1ml, 14.96mmol) to give the title compound (105mg) as a yellow
solid m.p. 117-118° δ H(d⁶ DMSO) 9.35 (1H, s), 8.83 (1H, d, \downarrow 2.2Hz),

8.38 (1H, d, J 5.3Hz), 8.13 (1H, dd, J 8.9, 2.4Hz), 7.27 (2H, s), 7.22 (1H, d, J 5.3Hz), 7.11 (1H, br t, J 5.0Hz), 3.77 (6H, s), 3.62 (3H, s), 3.29 (2H, q, J 6.3Hz), 2.71 (2H, t, J 6.3Hz) and 1.60 (2H, br s). MS m/z 397 (M+H)⁺.

5 **EXAMPLE 65**

4-(2-(2-Hydroxyethylamino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine

from the compound of Example 35 (300mg, 0.81mmol) and ethanolamine (1.0ml, 16.56mmol) to give the title compound (190mg) as an off-white solid m.p. 123°. $\delta\text{H}(\text{d}^6 \text{DMSO})$ 9.35 (1H, s), 8.82 (1H, d, J 2.2Hz), 8.38 (1H, d, J 5.3Hz), 8.12 (1H, dd, J 8.8, 2.4Hz), 7.26 (2H, s), 7.22 (1H, d, J 6.3Hz), 7.11 (1H, br t, J 5.0Hz), 6.59 (1H, d, J 8.8Hz), 4.73 (1H, br s), 3.77 (6H, s), 3.62 (3H, s), 3.57-3.52 (2H, m) and 3.45-3.38 (2H, m). MS m/z 398 (M+H)⁺.

15

EXAMPLE 66

4-(2-(3-Aminopropylamino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine

from the compound of Example 35 (250mg, 0.68mmol) and 1,3-diaminopropane (1.0ml, 11.89mmol) to give the title compound (120mg) as a yellow solid m.p. 152-153°. $\delta\text{H}(\text{d}^6 \text{DMSO})$ 9.34 (1H, s), 8.83 (1H, d, J 2.2Hz), 8.37 (1H, d, J 5.3Hz), 8.12 (1H, dd J 8.8, 2.4Hz), 7.26 (2H, s), 7.21 (1H, d, J 5.3Hz), 7.14 (1H, br t, J 5.5Hz), 6.53 (1H, d, J 8.8Hz), 3.77 (6H, s), 3.62 (3H, s), 3.34 (2H, q, J 7.0Hz), 2.61 (2H, t, J 6.7 Hz), 2.40 (2H, br s) and 1.65-1.56 (2H, m). MS m/z 411 (M+H)⁺.

25

EXAMPLE 67

4-(2-(2-Dimethylaminoethylamino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine

from the compound of Example 35 (250mg, 0.68mmol) and N,N-dimethylethylenediamine (0.75mol, 14.17mmol) to give the title compound (85mg) as an off-white solid m.p. 110-111°. MS m/z 425.5 (M+H)⁺.

30

EXAMPLE 68**4-(2-(4-Hydroxybutylamino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

from the compound of Example 35 (250mg, 0.68mmol) and 4-amino-1-butanol (0.750mg, 8.43mmol) to give the title compound (255mg) as a yellow solid m.p. 170-171°. MS m/z 4216 (M+H)⁺.

EXAMPLE 69

10 **4-(2-(2-(Pyrrolidin-1-yl)ethylamino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

from the compound of Example 35 (250mg, 0.68mmol) and N-(2-aminoethyl)pyrrolidine (0.75ml, 7.31mmol) to give the title compound (191mg) as a yellow solid m.p. 165-166°. MS m/z 451 (M+H)⁺.

15 **EXAMPLE 70**

4-(2-(2-Methylamino)ethyl(methyl)amino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine

from the compound of Example 35 (300mg, 0.81mmol) and N,N'-dimethylethylenediamine (0.61ml, 5.68mmol) to give the title compound (60mg) as a buff solid m.p. 115-116°. MS m/z 425 (M+H)⁺.

20 **EXAMPLE 71**

4-(2-(3-Isopropylaminopropylamino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine

25 from the compound of Example 35 (300mg, 0.81mmol) and N-isopropyl-1,3-propanediamine (0.51ml, 4.0mmol) to give the title compound (208mg) as a pale yellow solid m.p. 124-125°. MS m/z 453 (M+H)⁺.

30 **EXAMPLE 72**

4-(2-(1-Benzylpiperid-4-ylamino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine

from the compound of Example 35 (500mg, 1.35mmol) and 4-amino-1-benzylpiperidine (514mg, 2.7mmol) to give the title compound (370mg) as a yellow solid m.p. 113-114°. MS m/z 527 (M+H)⁺.

EXAMPLE 73**4-(2-(3-(Morpholino)propylamino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

5 from the compound of Example 35(300mg, 0.81mmol) and 3-(aminopropyl)morpholine (0.59ml, 4mmol) to give the title compound (300mg) as a white solid m.p. 165-166°. MS m/z 481 (M+H)⁺.

EXAMPLE 74**4-(2-(4-Methoxyphenylamino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

10 from the compound of Example 35 (300mg, 0.81mmol) and para-anisidine (200mg, 1.62mmol) to give the title compound (170mg) as a yellow solid m.p. 209-210°. MS m/z 459 (M+H)⁺.

EXAMPLE 75**4-(2-(3-Methylaminopropyl(methyl)amino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

15 In a manner analogous to Example 63 from the compound of Example 35 (500mg) and N,N'-dimethyl-1,3-propanediamine (1.62ml, 13.0mmol) to give the title compound (31mg) as a yellow solid. m.p. 103°. MS m/z 439 (M+H)⁺.

EXAMPLE 76**4-(2-(2-Hydroxycyclohexylamino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

25 from the compound of Example 35 (300mg, 0.81mmol), trans-2-amino-cyclohexanol hydrochloride (614mg, 4.05mmol) and triethylamine (0.56ml, 4.05mmol) to give the title compound (147mg) as a yellow solid m.p. 147-148°. MS m/z 452 (M+H)⁺.

30

EXAMPLE 77**4-(2-(3-Dimethylamino-2,2-dimethylpropylamino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

35 from the compound of Example 35 (300mg, 0.81mmol) and N,N,2,2-tetramethyl-1,3-propanediamine (527mg, 4.05mmol) to give the title compound (147mg) as a white solid m.p. 125°. MS m/z 467 (M+H)⁺.

EXAMPLE 78**4-(2-(3-Dimethylaminopropylamino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidine**

- 5 from the compound of Example 35 (300mg, 0.81mmol) and 3-dimethylaminopropylamine (248mg, 2.43mmol) to give the title compound (201mg) as a buff solid m.p. 170°. MS m/z 439 (M+H)⁺.

EXAMPLE 79

- 10 **4-(2-(2-Diethylaminoethyl(methyl)amino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidine**

from the compound of Example 35 (300mg, 0.81mmol) and N,N-diethyl-N'-methylethylenediamine (316mg, 2.43mmol) to give the title compound (372mg) as a buff solid m.p. 111°. MS m/z 467 (M+H)⁺.

15

EXAMPLE 80**4-(2-(3-Diethylaminopropylamino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

- 20 from the compound of Example 35 (300mg, 0.81mmol) and N,N-diethylaminopropylamine (282mg, 2.43mmol) to give the title compound (146mg) as a yellow solid m.p. 67°. MS m/z 453 (M+H)⁺.

EXAMPLE 81

- 25 **4-(2-(2-(1-Methylpyrrolidin-2-yl)ethylamino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

from the compound of Example 35 (300mg, 0.81mmol) and 2-(2-aminoethyl)-1-methylpyrrolidine (512mg, 4.0mmol) to give the title compound (190mg) as a yellow solid m.p. 176-177°. MS M/z 465 (M+H)⁺.

- 30 **EXAMPLE 82**

4-(2-(1-(Ethylpyrrolidin-2-yl)methylamino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine

- 35 from the compound of Example 35 (300mg), 0.81mmol) and 2-(amino-methyl)-1-ethylpyrrolidine (5.3mg, 4.0mmol) to give the title compound (240mg) as a yellow solid m.p. 81-82°. MS m/z 465 (M+H)⁺.

EXAMPLE 83**4-(2-(2-Dimethylaminoethyl(ethyl)amino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine.**

- from the compound of Example 35 (300mg, 0.81mmol) and N,N-dimethyl-N'-ethylethylenediamine (282mg, 2.43mmol) to give the title compound (193mg) as a yellow solid, m.p. 90°. MS m/z 453 (M+H)⁺.

EXAMPLE 84**4-(2-(4-Aminobutylamino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

- from the compound of Example 35 (300mg, 0.81mmol) and 1,4-butanediamine (705mg, 8.0mmol) as a yellow solid m.p. 209-201°. δ H(d⁶ DMSO) 9.35 (1H, s), 8.82 (1H, d, J 2.3Hz), 8.37 (1H, d, J 5.3Hz), 8.11 (1H, dd, J 8.9, 2.3Hz), 7.26 (2H, s), 7.21 (1H, d, J 5.3Hz), 7.15 (1H, br t, NH), 6.53 (1H, d, J 8.9Hz), 3.77 (6H, s), 3.61 (3H, s), 3.28 (2H, q, J 6.0Hz), 2.58 (2H, t, J 6.9Hz), 1.58-1.53 (2H, m) and 1.97-1.39 (2H, m). MS m/z 425 (M+H)⁺.

EXAMPLE 85

- 4-(2-(2-Diethylaminoethylamino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

from the compound of Example 35 (300mg, 0.81mmol) and N,N-diethylethylenediamine (282mg, 2.43mmol) to give the title compound (50mg) as a yellow solid m.p. 67°. MS m/z 453 (M+H)⁺.

25

EXAMPLE 86**4-(2-(3-(4-Methylpiperazin-1-yl)propylamino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

- from the compound of Example 35 (350mg, 0.94mmol) and 1-(3-aminopropyl)-4-methylpiperazine (629mg, 4.0mmol) to give the title compound (238mg) as a white solid m.p. 183-184°. MS m/z 494 (M+H)⁺.

30

Example 87**4-(2-(2-Dimethylaminoethyl(methyl)amino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)pyrimidine-2-amine**

from the compound of Example 35 (30mg, 0.81mmol) and N,N,N'-trimethylethylenediamine (400mg, 4.0mmol) to give the title compound (310g) as a yellow solid m.p. 97-98°. MS m/z 439 (M+H)⁺.

EXAMPLE 88

10 **4-(2-(Piperid-4-ylamino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine acetate**

To a solution of the compound of Example 72 (90mg, 0.15mmol) as the acetate salt in ethanol (10ml) was added cyclohexadiene (0.095ml, 1.0mmol) and 10% palladium on charcoal (100mg) and the reaction heated at reflux for 1h. On cooling the reaction was filtered through a pad of Celite® and the filtrate concentrated under reduced pressure to give the title compound (20mg) as a yellow solid m.p. 87-88°. MS m/z 437 (M+H)⁺.

EXAMPLE 89

20 **N4-Phenyl-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine**

A solution of N-phenyl-2-chloro-4-pyrimidineamine (1.83g, 8.91mmol) and 3,4,5-trimethoxyaniline (1.83g, 10.0mmol) in acetone (10ml) and water (15ml) containing concentrated hydrochloric acid (0.2ml) was heated at reflux for 10h. On cooling the acetone was removed under reduced pressure, and the reaction taken to pH10 with 4N NaOH and extracted with ethyl acetate (2x75ml). The combined organic layers were dried (MgSO₄), concentrated under reduced pressure and the residue crystallised from ethyl acetate to give the title compound (2.41g) as a colourless solid m.p. 191°. ¹H (CDCl₃) 8.04 (1H, d, J 5.8Hz), 7.37-7.34 (4H, m), 7.16-7.10 (1H, m), 7.05 (1H, br s), 6.85 (2H, s), 6.68 (1H, br s), 6.16 (1H, d, J 5.8Hz), 3.82 (3H, s) and 3.81 (6H, s). MS m/z 353 (M+H)⁺.

30 The pyrimidineamine used as starting material was prepared by heating a solution of 2,4-dichloropyrimidine (4.0g, 26.8mmol), aniline (2.46ml, 27.0mmol) and triethylamine (4.3ml, 30.0mmol) in ethanol (50ml) at reflux for 2h. On cooling a white precipitate was collected which was washed with cold ethanol (100ml) and recrystallised from ethyl acetate to give the

desired product (6.01g) as a colourless solid m.p. 186°. MS/mz 206 (M+H)⁺.

- 5 The compounds of Examples 90-110 were prepared in a similar manner to the compound of Example 89 using the starting material shown. In each case the pyrimidineamine starting material was prepared from the available compounds shown in a similar manner to the pyrimidineamine starting material of Example 89:

EXAMPLE 90

- 10 **N4-(4-(2-Hydroxyethoxy)phenyl)-N2-(3,4,5-trimethoxyphenyl)2,4-pyrimidineamine**

from 2-chloro-N-(4-(2-hydroxyethoxy)phenyl)-4-pyrimidineamine (187mg, 0.70mmol) and 3,4,5-trimethoxyaniline (146mg, 0.8mmol) to give the title compound (18mg) as a colourless solid m.p. 212°. δ H (d⁶ DMSO) 9.09 (1H, br s), 8.89 (1H, br s), 7.94 (1H, d, \downarrow 5.8Hz), 7.53 (2H, d, \downarrow 8.6Hz), 7.10 (2H, s), 6.86 (2H, d, \downarrow 8.6Hz), 6.10 (1H, d, \downarrow 5.8Hz), 4.83 (1H, t, \downarrow 5.6Hz), 3.95 (2H, t, \downarrow 4.9Hz), 3.71-3.68 (2H, m), 3.65 (6H, s) and 3.60 (3H,s). MS m/z 413 (M+H)⁺.

- 15 The pyrimidineamine starting material was prepared from 4-(2-hydroxyethoxy)aniline (481mg, 3.14mmol) 2,4-dichloropyrimidine (468mg, 3.14mmol) and triethylamine (0.46ml, 3.25mmol) to give the desired product as a yellow solid m.p. 169°. MS m/z 266(M+H)⁺.

EXAMPLE 91

- 25 **N4-(3,4-Methylenedioxyphenyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine**

from 2-chloro-N-(3,4-methylenedioxyphenyl)-4-pyrimidineamine (200mg, 0.85mmol) and 3,4,5-trimethoxyaniline (156mg, 0.85mmol) to give the title compound (187mg) as a buff solid m.p. 199°. MS m/z 397 (M+H)⁺.

- 30 The pyrimidineamine starting material was prepared from 3,4-methylenedioxyaniline (219mg, 1.60mmol) 2,4-dichloropyrimidine (250mg, 1.60mmol) and triethylamine (0.25ml, 1.80mmol) to give the desired product (215mg) as a buff solid m.p. 148°. MS m/z 250 (M+H)⁺.

EXAMPLE 92**N4-(3-Methoxyphenyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidine-diamine**

5 from 2-chloro-N4-(3-methoxyphenyl)-4-pyrimidineamine (195mg, 0.83mmol) and 3,4,5-trimethoxyaniline (152mg, 0.83mmol) to give the title compound (189mg) as a white solid m.p. 78°. MS m/z 383 (M+H)+.

The pyrimidineamine starting material was prepared from meta-anisidine (0.18mg, 1.6mmol), 2,4-dichloropyrimidine (250mg, 1.6mmol) and triethylamine (0.25mg, 1.8mmol) to give the desired product (200mg) as a
10 white solid m.p. 107°. MS m/z 236 (M+H)+.

EXAMPLE 93**N4-(3-(2-Hydroxyethyl)phenyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine**

15 from 2-chloro-N-(4-(2-hydroxyethyl)phenyl)-4-pyrimidineamine (250mg, 1.0mmol) and 3,4,5-trimethoxyaniline (184mg, 1.0mmol) to give the title compound (257mg) as a white solid m.p. 72°. MS m/z 397 (M+H)+.

The pyrimidineamine starting material was prepared from 2-(4-amino-phenyl)ethanol (219mg, 1.6mmol), 2,4-dichloropyrimidine (250mg, 1.6mmol) and triethylamine (0.25ml, 1.8mmol) to give the desired product
20 (289mg) as an orange solid m.p. 163°.

EXAMPLE 94**N4-(4-(Diethylamino)phenyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine**

25 from 2-chloro-N-(4-(diethylamino)phenyl)-4-pyrimidineamine (250mg, 0.9mmol) and 3,4,5-trimethoxyaniline (166mg, 0.9mmol) to give the title compound (80mg) as a green solid m.p. 87°. MS m/z 424.5 (M+H)+.

The pyrimidineamine starting material was prepared from N,N-diethyl-1,4-phenyldiamine (263mg, 1.6mmol), 2,4-dichloropyrimidine (250mg, 1.6mmol) and triethylamine (0.25ml, 1.8mmol) to give the desired prdocut
30 (410mg) as a green solid m.p. 212°. MS m/z 277 (M+H)+.

EXAMPLE 95**N4-(4-(3-Hydroxypropoxy)phenyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine**

5 from 2-chloro-N-(4-(3-hydroxypropoxy)phenyl)-4-pyrimidineamine (250mg, 0.89mmol) and trimethoxyaniline (164mg, 0.89mmol) to give the title compound (61mg) as a white solid m.p. 69°. MS m/z 427 (M+H)+.

The pyrimidineamine starting material was prepared (from 4-(3-hydroxypropoxy)aniline (267mg, 1.60mmol), 2,4-dichloropyrimidine (250mg, 1.60mmol) and triethylamine (0.25ml, 1.80mmol) to give the desired
10 product (383mg) as a pale orange solid m.p. 181°. MS m/z 280 (M+H)+.

EXAMPLE 96**N4-(3,4-Dimethoxyphenyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine**

15 from 2-chloro-N-(3,4-dimethoxyphenyl)-4-pyrimidineamine (250mg) and 3,4,5-trimethoxyaniline (189mg, 1.0mmol) to give the title compound (160mg) as a light pink solid m.p. 86°. MS m/z 413 (M+H)+.

The pyrimidineamine starting material was prepared from aminoveratrole (257mg, 1.60mmol), 2,4-dichloropyrimidine (2.50mg, 1.60mmol) and
20 triethylamine (0.25ml, 1.80mmol) to give the desired product (357mg) as a purple solid m.p. 104° MS m/z 266 (M+H)+.

EXAMPLE 97

25 **N4-(4-(4-Methylpiperazin-1-yl)phenyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine**

from 2-chloro-N-(4-(4-methylpiperazin-1-yl)phenyl)-4-pyrimidineamine (110mg, 0.40mmol) and 3,4,5-trimethoxyaniline (66mg, 0.40mmol) to give the title compound (80mg) as a white solid m.p. 104°. MS m/z 451 (M+H)+.

30 The pyrimidineamine starting material was prepared from 4-(4-methylpiperazin-1-yl)aniline (511mg, 2.67mmol), 2,4-dichloropyrimidine (398mg, 2.67 mmol) and triethylamine (10.4ml, 2.94mmol) to give the desired product (110mg) as a white solid m.p. 122°.

EXAMPLE 98**N4-(3,5-Dimethoxyphenyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine**

5 from 2-chloro-N-(3,5-dimethoxyphenyl)-4-pyrimidineamine (150mg, 0.56mmol) and 3,4,5-trimethoxyaniline (104mg, 0.56mmol) to give the title compound (162mg) as a white solid m.p. 180°. MS m/z 413 (M+H)⁺.

The pyrimidineamine starting material was prepared from 3,5-dimethoxyaniline (257mg, 1.60mmol), 2,4-dichloropyrimidine (250mg, 1.60mmol) and triethylamine (0.25ml, 1.80mmol) to give the desired product (237mg) 10 as a white solid m.p. 225°. MS m/z 266 (M+H)⁺.

EXAMPLE 99**N4-(4-Morpholinophenyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidine-diamine**

15 from 2-chloro-N-(4-morpholinophenyl)-4-pyrimidineamine (300mg, 1.03mmol) and 3,4,5-trimethoxyaniline (189mg, 1.03mmol) to give the title compound (230mg) as a white solid m.p. 214°. MS m/z 438 (M+H)⁺.

The pyrimidineamine starting material was prepared from 4-morpholinoaniline (285mg, 1.60mmol), 2,4-dichloropyrimidine (205mg, 1.60mmol) and triethylamine (0.25ml, 1.80mmol) to give the desired product (322mg) 20 as a white solid m.p. 221°. MS m/z 291 (M+H)⁺.

EXAMPLE 100**N4-(3-Trifluoromethoxyphenyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine**

25 from 2-chloro-N-(3-trifluoromethoxyphenyl)-4-pyrimidineamine (230mg, 0.79mmol) and 3,4,5-trimethoxyaniline (145mg, 0.79mmol) to give the title compound (220mg) as a white solid m.p. 154-156°. MS m/z 437 (M+H)⁺.

The pyrimidineamine starting material was prepared from 3-trifluoromethoxyaniline (354mg, 2.0mmol) 2,4-dichloropyrimidine (218mg, 2.0mmol) and triethylamine (0.3ml, 2.20mmol) to give the desired product (240mg) 30 as a white solid m.p. 161-163°. MS m/z 290 (M+H)⁺.

EXAMPLE 101**N4-(2-Methylbenzimidazol-5-yl)-N2-(3,4,5-trimethoxyphenylamino)-2,4-pyrimidinediamine**

5 from 2-chloro-N-(2-methylbenzimidazol-5-yl)-4-pyrimidineamine (250mg, 0.96mmol) and 3,4,5-trimethoxyaniline (176mg, 0.96mmol) to give the title compound (36mg) as a white solid m.p. 179°. MS m/z 407 (M+H)⁺.

The pyrimidineamine starting material was prepared from 5-amino-2-methylbenzimidazole (2147mg, 1.68mmol), 2,4-dichloropyrimidine (250mg, 1.68mmol) and triethylamine (0.25ml, 1.80mmol) to give the
10 desired product (344mg) as a pale pink solid m.p. >300°. MS m/z 260 (M+H)⁺.

EXAMPLE 102

15 **N4-(4-(Dimethylamino)phenyl)-N2-(3,4,5-trimethoxyphenylamino)-2,4-pyrimidinediamine**

from 2-chloro-N-(4-(dimethylamino)phenyl)-4-pyrimidineamine (250mg, 1.0mmol) and 2,3,4-trimethoxyaniline (184mg, 1.0mmol) to give the title compound (127mg) as a grey solid m.p. 215°. MS m/z 396 (M+H)⁺.

The pyrimidineamine starting material was prepared from N,N-dimethyl-
20 1,4-phenylenediamine (228mg, 1.68mmol) 2,4-dichloropyrimidine (250mg, 1.68mmol) and triethylamine (0.25ml, 1.80mmol) to give the desired product (402mg) as a white solid m.p. 212°.

EXAMPLE 103

25 **N4-(4-(Ethoxycarbonylmethyl)phenyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine**

from 2-chloro-N-(4-ethoxycarbonylmethyl)phenyl)-4-pyrimidineamine (1.74g, 5.97mmol) and 3,4,5-trimethoxyphenylaniline (1.09g, 5.97mmol) to give the title compound (1.01g) as a white solid m.p. 143-144°. δ H
30 (CDCl₃) 8.04 (1H, d, J 5.8Hz), 7.32 (2H, d, J 8.6Hz), 7.26 (2H, d, J 8.6Hz), 7.01 (1H, br s), 6.86 (2H, s), 6.64 (1H, br s), 6.15 (1H, d, J 5.8Hz), 4.17 (2H, q, J 7.1Hz), 3.82² (3H, s), 3.81 (6H, s), 3.60 (2H, s) and 1.27 (3H, t, J 7.1Hz). MS m/z 439 (M+H)⁺

The pyrimidineamine starting material was prepared from ethyl 4-amino-
35 phenylacetate (1.79g, 10.0mmol) 2,4-dichloropyrimidine (1.49g,

10.0mmol) and triethylamine (2.8ml, 20mmol) to give the desired product (2.88g) as a white solid, m.p. 151-152°.

EXAMPLE 104

5 **N4-(4-Benzyloxyphenyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidine-diamine**

from N-(4-benzyloxyphenyl)-2-chloro-4-pyrimidineamine (25.0g, 80.0mmol) and 3,4,5-trimethoxyaniline (14.7g, 80.0mmol) to give the title compound (18.6g) as a white solid m.p. 161-164°. MS m/z 459 (M+H)+.

10 The pyrimidineamine starting material was prepared from 4-benzyloxyaniline (20.0g, 0.1mol), 2,4-dichloropyrimidine (15.0g, 0.1mol) and triethylamine (15.0ml, 0.11mol) to give the desired product (30.0g) as a cream solid m.p. 198°. MS m/z 312 (M+H)+.

15 **EXAMPLE 105**

N4-(3-Ethoxyphenyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidine-diamine

from 2-chloro-N-(3-ethoxyphenyl)-4-pyrimidineamine (125mg, 0.5mmol) and 3,4,5-trimethoxyaniline (92mg, 0.5mmol) to give the title compound (157mg) as a white solid m.p. 79°. MS m/z 397 (M+H)+.

20 The pyrimidineamine starting material was prepared from meta-phenetidine (0.15ml, 1.68mmol), 2,4-dichloropyrimidine (250mg, 1.68mmol) and triethylamine (0.25ml, 1.80mmol) to give the desired product (125mg) as a cream solid m.p. 97°.

25

EXAMPLE 106

N4-(4-Benzyloxycarbonylmethylphenyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine

30 from N-(4-(benzyloxy-carbonylmethyl)phenyl)-2-chloro-4-pyrimidineamine (3.14g, 8.90mmol) and 3,4,5-trimethoxyaniline (1.63g, 8.90mmol) to give the title compound (3.47g) as a white solid m.p. 141-143°. δ H (CDCl₃) 8.04 (1H, d, J 5.8Hz), 7.36-7.24 (9H, m), 7.15 (1H, br s), 6.86 (2H, s), 6.74 (1H, br s), 6.15 (1H, d, J 5.8Hz), 5.15 (2H, s), 3.82 (3H, s), 3.81 (6H, s) and 3.66 (2H, s). MS m/z 501 (M+H)+.

35 The pyrimidineamine starting material was prepared from benzyl 4-aminophenylacetate (4.82g, 20.0mmol), 2,4-dichloropyrimidine (2.98g,

20.0mmol) and triethylamine (5.60ml, 40.0mmol) to give the desired product (3.24g) as a white solid m.p. 149°. MS m/z 354 (M+H).⁺

EXAMPLE 107

5 **N4-(4-Methoxyphenyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidine-diamine**

from 2-chloro-N-(4-methoxyphenyl)-4-pyrimidineamine (315mg, 1.30mmol) and 3,4,5-trimethoxyaniline (270mg, 1.50mmol) to give the title compound (470mg) as a white solid m.p. 210°. MS m/z 383 (M+H).⁺

- 10 The pyrimidineamine starting material was prepared from para-anisidine (616mg, 5.0mmol), 2,4-dichloropyrimidine (745mg, 5.0mmol) and triethylamine (0.77ml, 5.5mmol) to give the desired compound (450mg) as a white solid m.p. 250°.

15 **EXAMPLE 108**

N4-(3-Benzyloxyphenyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidine-amine

from N-(3-benzyloxyphenyl)-2-chloro-4-pyrimidineamine (2.0g, 6.4mmol) and 3,4,5-trimethoxyaniline (1.18g, 6.4mmol) to give the title compound (2.60g) as a white solid m.p. 186°. MS m/z 459 (M+H).⁺

- 20 The pyrimidineamine starting material was prepared from 3-benzyloxyaniline (5.0g, 25.0mmol), 2,4-dichloropyrimidine (3.7g, 25.0mmol) and triethylamine (3.8ml, 27.5mmol) to give the desired compound (4.95g) as a white solid m.p. 119°. MS m/z 312 (M+H).⁺

25

EXAMPLE 109

N4-(1-Phenylsulphonylindol-5-yl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine

from 2-chloro-N-(1-phenylsulphonylindol-5-yl)-4-pyrimidineamine (1.09g, 2.70mmol) and 3,4,5-trimethoxyaniline (496mg, 2.70mmol) to give the title compound (950mg) as a white solid m.p. 127°. MS m/z 532 (M+H).⁺

- 30 The pyrimidineamine starting material was prepared from 5-amino-1-phenylsulphonylindole (1.0g, 3.9mmol), 2,4-dichloropyrimidine (577mg, 3.9mmol) and triethylamine (0.61ml) to give the desired product (1.20g) as
35 an orange solid m.p. 158°. MS m/z 385 (M+H).⁺

EXAMPLE 110**N4-Cyclohexyl-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine**

from 2-chloro-N-cyclohexylamino-4-pyrimidineamine (200mg, 0.9mmol) and 3,4,5-trimethoxyaniline (173mg, 0.9mmol) to give the title compound (173mg) as a white solid m.p. 160-161°. δ H (CDCl₃) 7.91 (1H, d, \downarrow 5.9Hz), 6.90 (1H, s), 6.87 (2H, s), 5.81 (1H, d, \downarrow 5.9Hz), 4.63 (1H, br s), 3.86 (6H, s), 3.81 (3H, s), 2.00 (2H, m), 1.70 (4H, m) and 1.25 (4H, m). MS m/z 359 (M+H)⁺

The pyrimidineamine starting material was prepared from cyclohexylamine (0.4ml, 3.4mmol), 2,4-dichloropyrimidine (500mg, 3.4mmol) and triethylamine (0.5ml, 3.7mmol) to give the desired product (270mg) as a white solid m.p. 128°. MS m/z 212 (M+H)⁺.

EXAMPLE 111**N4-(1-Benzylpiperid-4-yl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidine-diamine**

from N-(1-benzylpiperid-4-yl)-2-chloro-4-pyrimidineamine (390mg, 1.3mmol) and 3,4,5-trimethoxyaniline (236mg, 1.3mmol) to give the title compound (126mg) as a white solid m.p. 147°. δ H (CDCl₃) 7.90 (1H, d, \downarrow 5.8Hz), 7.28 (5H, m), 6.94 (1H, s), 6.86 (2H, s), 5.81 (1H, d, \downarrow 5.8Hz), 4.62 (1H, br s), 3.85 (6H, s), 3.81 (3H, s), 3.53 (2H, s), 2.83 (2H, m), 2.16 (2H, m), 2.01 (2H, m) and 1.52 (2H, m).

The pyrimidineamine starting material was prepared from 4-amino-1-benzylpiperidine (0.7ml, 3.4mmol), 2,4-dichloropyrimidine (500mg, 3.4mmol) and triethylamine (0.5ml, 3.7mmol) to give the desired product (650mg) as a white solid m.p. 136°. MS m/z 303 (M+H)⁺.

EXAMPLE 112**N4-Benzyl-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine**

from N-benzyl-2-chloro-4-pyrimidineamine (1.5g, 6.83mmol) and 3,4,5-trimethoxyaniline (1.31g, 7.17mmol) to give the title compound (2.30g) as a white solid m.p. 167-168°. MS m/z 367 (M+H)⁺.

The pyrimidineamine starting material was prepared from benzylamine (3.67ml, 33.56mmol), 2,4-dichloropyrimidine (5.0g, 33.56mmol) and triethylamine (5.14ml, 36.9mmol) to give the desired product (4.21g) as a white solid. m.p. 135-136°. MS m/z 220 (M+H)⁺.

EXAMPLE 113**N2-(3,4-Dimethoxy-5-(methylthio)phenyl)-N4-(3-methoxyphenyl)-2,4-pyrimidinediamine**

5 from 2-chloro-N-(3-methoxy-phenyl)-4-pyrimidineamine (0.59g, 2.51mmol) [see Example 92] and 3,4-dimethoxy-5-(methylthio)aniline (0.50g, 2.51mmol) to give the title compound (0.91g) as a white solid m.p. 68-70°. δ H (CDCl₃) 8.02 (1H, d, \downarrow 5.8Hz), 7.40 (1H, br, s), 7.26 (1H, m), 7.08 (1H, d, \downarrow 2.2Hz), 6.90 (4H, m), 6.68 (1H, dd, \downarrow 8.2, 2.2Hz), 6.20 (1H, d, \downarrow 5.8Hz), 3.82 (3H, s), 3.79 (3H, s), 3.77 (3H, s) and 2.35 (3H, s), MS m/z 399 (M+H)⁺.

15 The aniline starting material was prepared by treating a solution of 1,2-dimethoxy-3-(methylsulphonyl)-5-nitrobenzene (1.64g, 6.69mmol) in methanol (20ml) and concentrated hydrochloric acid (20ml) was treated with anhydrous tin (II) chloride (7.15g, 37.7mmol) and the resulting mixture refluxed for 1.25h. On cooling to room temperature the mixture was poured into excess 1M NaOH solution and extracted with CH₂Cl₂. The organic extract was dried (MgSO₄) and evaporated to afford the desired product (1.33g) as an off-white solid m.p. 106-107°. MS m/z 199 (M+H)⁺.

20

EXAMPLE 114**N2-(3,4-Dimethoxy-5-methylphenyl)-N4-(3-methoxyphenyl)-2,4-pyrimidinediamine**

25 from 2-chloro-N-(3-methoxyphenyl)-4-pyrimidineamine (1.93g) [see Example 92] and 3,4-dimethoxy-5-methylaniline (1.50g) to give the title compound (2.53g) as a pale pink solid m.p. 66-68°. δ H (CDCl₃) 8.02 (1H, d, \downarrow 5.8Hz), 7.39 (1H, br s), 7.23 (1H, t, \downarrow 8.1Hz), 7.08 (1H, m), 7.05 (1H, br s), 6.97-6.91 (2H, m), 6.86 (1H, d, \downarrow 2.0Hz), 6.69-6.65 (1H, m), 6.18 (1H, d, \downarrow 5.8Hz), 3.77 (3H, s), 3.75 (6H, s) and 2.22 (3H, s). MS m/z 367 (M+H)⁺.

30

The aniline used as starting material was prepared according to the methods of I Sanchez *et al*: Tetrahedron **41**, 2355 (1985).

EXAMPLE 115**N2-(3,4-Dimethoxyphenyl)-N4-(3-methoxyphenyl)-2,4-pyrimidinediamine**

- from 2-chloro-N-(3-methoxyphenyl)-4-pyrimidineamine (0.77g, 3.27mmol)
5 [see Example 92] and 3,4-dimethoxyaniline (0.50g, 3.27mmol) to give the
title compound (0.89g) as a beige solid m.p. 135-138°. δ H (CDCl₃) 8.03
(1H, d, \downarrow 5.8Hz), 7.22 (2H, m), 7.13 (1H, br s), 7.02 (1H, dd, \downarrow 8.6, 2.3Hz),
6.97 (1H, m), 6.92 (1H, m), 6.82 (1H, d, \downarrow 87.6Hz), 6.77 (1H, br s), 6.68
(1H, dd, \downarrow 8.2, 2.3Hz), 6.18 (1H, d, \downarrow 5.8Hz), 3.86 (3H, s), 3.83 (3H, s) and
10 3.77 (3H, s). MS m/z 353 (M+H)⁺.

EXAMPLE 116**N2-(3-Chloro-4,5-dimethoxyphenyl)-N4-(3-methoxyphenyl)-2,4-pyrimidinediamine hydrochloride**

- 15 from 2-chloro-N-(3-methoxyphenyl)-4-pyrimidineamine (0.50g, 2.13mmol)
[see Example 92] and 3-chloro-4,5-dimethoxyaniline (0.40g, 2.13mmol) to
give the title compound (0.55g) as a white solid m.p. 204-205°. δ H (d⁶
DMSO) 11.19 (1H, br s), 10.85 (1H, br s), 8.00 (1H, d, \downarrow 7.1Hz), 7.23 (4H,
m), 7.12 (1H, d, \downarrow 4.4Hz), 6.75 (1H, m), 6.59 (1H, d, \downarrow 7.1Hz), 3.74 (3H, s),
20 3.69 (3H, s) and 3.65 (3H, s). MS m/z 387 (M+H)⁺.

The aniline starting material was prepared in a similar manner to the
analogous aniline of Example 113, from 1-chloro-2,3-dimethoxy-5-
nitrobenzene (1.08g, 5.82mmol), to give the desired product (0.85g) as a
white solid m.p. 66-68°. MS m/z 188 (M+H)⁺.

- 25 The 1-chloro-2,3-dimethoxy-5-nitrobenzene was prepared by heating a
solution of 3-chloro-4,5-dimethoxybenzoic acid (3.50g, 16.2mmol) in
glacial acetic acid (15ml) and 70% nitric acid (15ml) at 60° for 1h. The
reaction was poured onto ice-water and the white precipitate which formed
was filtered off, washed with water and dried *in vacuo* and washed
30 thoroughly with hexane. The hexane washings were evaporated and the
residue subjected to column chromatography [silica 20% ethyl acetate-
hexane] to give the desired product (1.03g) as a white solid m.p. 104-
105°. MS m/z 217 (M+H)⁺.

- The acid used as starting material was prepared according to the method
35 of Y. Ohtani *et al.* Acta. Chem. Scand., Ser B. **B36**, 613 (1982).

EXAMPLE 117**N2-(3-Benzoyloxy-4,5-dimethoxyphenyl)-N4-(3-methoxyphenyl)-2,4-pyrimidinediamine**

from 2-chloro-N-(3-methoxyphenyl)-4-pyrimidineamine (162mg, 0.69mmol)
5 [see Example 92] and 3-benzoyloxy-4,5-dimethoxyaniline (180mg, 0.69mmol) to afford the title compound (255mg) as an off-white solid m.p. 83-84°. δ H (CDCl₃) 8.01 (1H, d, \downarrow 5.8Hz), 7.45-7.22 (6H, m), 6.95 (3H, m), 6.90 (1H, d, \downarrow 7.6Hz), 6.79 (1H, d, \downarrow 2.4Hz), 6.68 (1H, d, \downarrow 7.6Hz), 6.48 (1H, s), 6.17 (1H, d, \downarrow 5.8Hz), 5.11 (2H, s), 3.86 (3H, s), 3.82 (3H, s) and
10 3.78 (3H, s). MS m/z 369 (M+H)⁺.

The 3-benzoyloxy-4,5-dimethoxyaniline was prepared by heating a solution of 1-benzoyloxy-2,3-dimethoxy-5-nitrobenzene (1.80g, 6.23mmol) in ethanol (15ml) with saturated aqueous sodium hydrosulphite (20ml) at reflux for 2h. An additional quantity of sodium hydrosulphite (20ml) was
15 added and reflux continued for a further 4h. The reaction mixture was reduced to a small volume then diluted with water and extracted three times with ethyl acetate. The organic phase was dried (MgSO₄) and evaporated to give the crude product which was recrystallised from ether-hexane to give the desired product (356mg) as a white crystalline solid
20 m.p. 118-120°. MS m/z 260 (M+H)⁺.

The 1-benzoyloxy-2,3-dimethoxy-5-nitrobenzene used as starting material was prepared by treating a solution of 1-benzoyl-2,3-dimethoxybenzene (7.25g, 29.66mmol) in glacial acetic acid (15ml) portionwise with 70% nitric acid (2.84ml) at room temperature. After 2h, the reaction mixture was
25 poured into ice-water and extracted with ethyl acetate. The organic phase was washed with 1M NaOH, then brine, dried (MgSO₄) and evaporated. The crude product was purified by column chromatography [silica 20% ethyl acetate-hexane] to give, after recrystallisation from CH₂Cl₂-hexane, the desired product (1.89g) as a white solid m.p. 101-102°. MS m/z
30 (relative intensity) 289 [(M+H)⁺, 2%] 91 [100%].

1-Benzoyloxy-2,3-dimethoxybenzene was prepared by the method of F Dietl *et al* Synthesis, 626 (1985).

EXAMPLE 118**N2-(3-Acetamido-4,5-dimethoxyphenyl)-N4-(3-methoxyphenyl)2,4-pyrimidinediamine**

- from 2-chloro-N-(3-methoxyphenyl)-4-pyrimidineamine (0.81g, 3.43mmol) [see Example 92] and 3-acetamido-4,5-dimethoxyaniline (0.70g, 3.43mmol) to give the title compound (573mg) as an off-white solid m.p. 195-197°. δ H (d^6 DMSO) 9.26 (1H, s), 9.10 (1H, s), 9.01 (1H, s), 7.97 (1H, d, J 5.7Hz), 7.74 (1H, s), 7.46-7.25 (3H, m), 7.15 (1H, t, J 8.1Hz), 6.54 (1H, d, J 8.1Hz), 6.18 (1H, d, J 5.7Hz), 3.67 (9H, m) and 2.06 (3H, s). MS m/z 410 (M+H)⁺.
- The 3-acetamido-4,5-dimethoxyaniline was prepared in an analogous manner to the compound of Example 38, from 1-acetamido-2,3-dimethoxy-5-nitrobenzene (1.65g, 6.88mmol) to afford the desired product (1.37g) as a brown solid m.p. 158-160°. MS m/z 211 (M+H)⁺.
- The 1-acetamido-2,3-dimethoxy-5-nitrobenzene was prepared in an analogous manner to the nitrobenzene of Example 117 from 1-acetamido-2,3-dimethoxybenzene (1.91g, 9.8mmol) to give the desired product (1.70g) as a white solid m.p. 83°. MS m/z 241 (M+H)⁺.
- The 1-acetamido-2,3-dimethoxybenzene was prepared by treating a solution of 2,3-dimethoxyaniline (1.70g, 11.1mmol) in carbon tetrachloride (25ml) with methanol (0.2ml) followed by acetic anhydride (3.67g, 36.0mmol) and heating the resulting mixture at reflux for 1h. The cooled reaction mixture was diluted with CH₂Cl₂, washed with water, dried (MgSO₄) and evaporated to give the desired product (2.11g) as a beige solid m.p. 164-165°. MS m/z 196 (M+H)⁺.

EXAMPLE 119**5-Bromo-N2,N4-Bis(trimethoxyphenyl)-2,4-pyrimidinediamine**

- from 5-bromo-2-chloro-N-(3,4,5-trimethoxyphenyl)-4-pyrimidineamine (936mg, 2.5mmol) and 3,4,5-trimethoxyaniline (459mg, 2.5mmol) to give the title compound (160mg) as a white solid m.p. 186-191°. δ H (CDCl₃) 8.16 (1H, s), 7.45 (1H, br s), 6.98 (1H, br s), 6.77 (2H, s), 6.74 (2H, s), 3.83 (3H, s), 3.80 (3H, s), 3.71 (6H, s), 3.66 (6H, s), MS m/z 523 [(M+H)⁺, ⁸¹Br, 100%] 521 [(M+H)⁺, ⁷⁹Br, 100%].
- The pyrimidineamine starting material was prepared from 5-bromo-2,4-dichloropyrimidine (1.14g, 5.0mmol), 3,4,5-trimethoxyaniline (916mg,

5.0mmol) and triethylamine (0.77ml, 5.5mmol) following the method used for the pyrimidineamine starting material in Example 89. This gave the desired compound as a white solid (1.71g) m.p. 185-186°. MS m/z 378 [(M+H)⁺, 22%], 376 [(M+H)⁺, 100%], 374 [(M+H)⁺, 78%].

5

EXAMPLE 120**N4-(4-Hydroxyphenyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidine-diamine**

10 In a manner analogous to the preparation of the compound of Example 49 from the compound of Example 104 (3.0g, 6.55mmol), ammonium formate (1.20g, 19.60mmol) and 10% palladium on carbon (300mg) to give the title compound (2.40g) as a white solid m.p. 112-115°. δ H (CDCl₃) 7/96)1H, d, \downarrow 5.9Hz), 7.16 (2H, d, \downarrow 8.7Hz), 7.06 (1H, s), 6.83 (1H, s), 6.80 (3H, m), 6.68 (1H, s), 6.01 (1H, d, \downarrow 5.9Hz), 3.79 (3H, s) and 3.78 (6H, s). MS m/z 15 369 (M+H)⁺.

EXAMPLE 121**N4-(4-(3-Aminopropoxy)phenyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine dihydrochloride**

20 In a manner analogous to the preparation of the compound of Example 48 from N4-(4-(3-phthalimidopropoxy)phenyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine (400mg, 0.72mmol) and hydrazine monohydrate (0.1ml, 2.16mmol) to give the title compound (108mg) as a buff solid m.p. 258°. δ H (d⁶ DMSO) 11.04 (1H, s), 10.54 (1H, s), 8.08 (3H, s), 7.91 (1H, d, \downarrow 7.3Hz), 7.55 (2H, m), 6.86 (2H, d, \downarrow 8.7Hz), 6.78 (2H, s), 6.49 (1H, s), 25 4.03 (2H, m), 3.65 (9H, s), 2.94 (2H, m) and 2.02 (2H, m). MS m/z 426 (M+H)⁺.

30 The pyrimidinediimine starting material was prepared according to the method of Example 45 from the compound of Example 120 and 3-bromopropylphthalimide

EXAMPLE 122**N4-(4-(Carboxymethyl)phenyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine**

35 The compound of Example 103 (250mg, 0.57mmol) in a solution of ethanol (10ml) containing 6N aqueous NaOH (2.0ml) was heated a reflux

for 2h. On cooling the ethanol was removed under reduced pressure and the basic solution brought to pH 5 with 2M hydrochloric acid. The resulting solution was concentrated under reduced pressure, and the residue taken up in hot ethanol (20ml). This solution was filtered and the filtrate
5 evaporated under reduced pressure to give a residue which was crystallised from ethanol to give the title compound (89mg) as a white solid m.p. 144-150°. MS m/z 411 (M+H)⁺.

EXAMPLE 123

10 **N4-(4-(*tert*-Butoxycarbonylmethoxy)phenyl)-N2-(3,4,5-trimethoxy-phenyl)-2,4-pyrimidinediamine**

In a manner analogous to the preparation of the compound of Example 50, from the compound of Example 120 (500mg, 1.40mmol), *tert*-butyl bromoacetate (0.2ml, 1.40mmol) and NaH (60% dispersion in oil) (60ml, 1.50mmol) to give the title compound (380mg) as a green solid m.p. 139°. MS m/z 483 (M+H)⁺.
15

EXAMPLE 124

20 **N4-(4-(2-Hydroxyethyl)phenyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine**

In a manner analogous to the preparation of the compound of Example 62 (267mg, 0.5mmol) from the compound of Example 106 and LiAlH₄ (40mg, 1.1mmol) to give the title compound (108mg) as a light pink solid m.p. 172°. MS m/z 397 (M+H)⁺.
25

EXAMPLE 125

30 **N4-(4-(Carboxymethoxy)phenyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine**

A solution of the compound of Example 123 (300mg, 0.62mmol) in CH₂Cl₂ (20ml) was treated with trifluoroacetic acid (3.0ml) and stirred at room temperature for 5h. After this time the reaction was concentrated under reduced pressure to give a brown oil, which was subjected to column chromatography [silica CH₂Cl₂/methanol/acetic acid/water 86.5:10:2:1.5] to give the title compound (123mg) as a white solid m.p. 159° after
35 recrystallisation from methanol. MS m/z 427 (M+H)⁺.

EXAMPLE 126**4-Phenoxy-N2-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

2-Methylsulphonyl-4-phenoxy-pyrimidine (0.76g, 3.04mmol) and 3,4,5-trimethoxyaniline (0.56g, 4.0mmol) were heated as a melt at 140° for 2h.

- 5 On cooling the residue was partitioned between ethyl acetate (50ml) and 2M hydrochloric acid (50ml), and the organic layer was washed with water (50ml), dried (MgSO₄) and concentrated under reduced pressure. The residue was subjected to column chromatography [silica 20% ethyl acetate-hexane] to give the title compound (18mg) as a white solid m.p. 149° after recrystallisation from ethyl acetate. δ H (CDCl₃) 8.26 (1H, d, \downarrow 5.6Hz), 7.39 (2H, m), 7.26 (1H, m), 7.18-7.15 (2H, m), 6.73 (2H, s), 6.31 (1H, d, \downarrow 5.6Hz), 3.77 (3H, s) and 3.64 (6H, s). MS m/z 354 (M+H)⁺.

- 10 The pyrimidine used as starting material was prepared by the addition of 50% 3-chloroperoxybenzoic acid (15.19g, 88.0mmol) to a solution of 4-phenoxy-2-thiomethylpyrimidine (5.0g, 22.9mmol) in CH₂Cl₂ (100ml) at 0°. The resulting mixture was allowed to rise to room temperature over 12h, and was then washed with 2M aqueous NaOH (2x100ml), saturated aqueous sodium sulphite (2x100ml), dried (MgSO₄) and concentrated under reduced pressure. The residue was subjected to column chromatography to give the desired product (0.81g) as a colourless oil. δ H (CDCl₃) 8.49 (1H, d, \downarrow 5.5Hz), 7.45 (2H, t, \downarrow 8.0Hz), 7.33-7.26 (1H, m), 7.19-7.16 (2H, m), 6.85 (1H, d, \downarrow 5.5Hz) and 3.23 (3H, s).

- 20 The 4-phenoxy-2-thiomethylpyrimidine used as starting material was prepared by heating a solution of 4-fluoro-2-thiomethylpyridine [N. Plé *et al.*, J. Het. Chem. 31, 1311-1315 (1994)] (9.98g, 69.3mmol) and phenol (6.58g, 70.0mmol) in dry DMF (100ml) in the presence of caesium carbonate (22.8g, 70.0mmol) at 40° for 6h. The reaction was concentrated *in vacuo* and the residue was partitioned between ethyl acetate (100ml) and water (100ml). The organic layer was dried (MgSO₄), concentrated under reduced pressure and the residue subjected to column chromatography [silica 40% ethyl acetate-hexane] to give the desired product (7.01g) as a colourless oil. δ H (CDCl₃) 8.33 (1H, dd, \downarrow 4.7, 0.9Hz), 7.40 (2H, dd, 8.0, 1.1Hz), 7.26 (1H, m), 7.23 (2H, m), 6.46 (1H, dd, \downarrow 4.7, 0.9Hz) and 2.37 (3H, s). MS m/z 219 (M+H)⁺.

EXAMPLE 127**N,N'-Bis(3,4,5-Trimethoxyphenyl)-2,4-pyrimidinediamine**

A solution of 2,4-dichloropyrimidine (1.0g, 6.7mmol), 3,4,5-trimethoxyaniline (2.5g, 13.4mmol) and triethylamine (1.8ml, 12.6mmol) was heated at reflux for 5h. On cooling the resulting precipitate was collected and recrystallised from ethyl acetate to give the title compound (450mg) as a purple solid m.p. 180°. MS m/z 443 (M+H)⁺.

EXAMPLE 128**10 4-Benzoyl-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

To a solution of N-methoxy-N-methyl-2-(3,4,5-trimethoxyphenylamino)-pyrimidine-4-carboxamide (0.82g, 2.30mmol) in THF (40ml) cooled to -78° under a nitrogen atmosphere was added phenylmagnesium bromide [Aldrich 1M/THF] (6.90ml, 6.90mmol) dropwise. On completion of addition the reaction was left to warm to room temperature over 3h and aqueous NH₄Cl solution (10ml) was then added and the mixture concentrated under reduced pressure. The residue was partitioned between water (20ml) and ethyl acetate (40ml), the organic layer dried (MgSO₄) and evaporated. The residue was subjected to column chromatography [silica 50% hexane-ethyl acetate] and after recrystallisation from ethyl acetate-hexane the title compound (123mg) was obtained as a yellow solid m.p 135°. δ H (CDCl₃) 8.65 (1H, d, \downarrow 4.8Hz), 8.05 (2H, dd, \downarrow 5.1, 2.0Hz), 7.61 (1H, dd, \downarrow 5.3, 2.0Hz), 7.46 (2H, m), 7.24 (1H, d, \downarrow 4.8Hz), 3.79 (3H, s) and 3.63 (6H, s). MS m/z 366 (M+H)⁺.

25 The pyrimidine-4-carboxamide used as starting material was prepared by treating a solution of 4-carboxy-N-(3,4,5-trimethoxyphenyl)-2-pyrimidine-amine hydrochloride (2.0g, 5.86mmol) and triethylamine (2.0ml, 14.0mmol) in CH₂Cl₂ (40ml), cooled to -20° under a nitrogen atmosphere, with isobutylchloroformate (1.1ml, 7.0mmol). The resulting mixture was stirred for 20min. before N,O-dimethylhydroxylamine hydrochloride (683mg, 7.0mmol) and triethylamine (1.0ml, 7.0mmol) were added and the reaction allowed to warm to room temperature over 3h. After washing with 2M hydrochloric acid (1x50ml) and 2M NaOH solution (1x50ml), the reaction was dried (MgSO₄) and concentrated under reduced pressure.

30 The residue was subjected to column chromatography [silica ethyl acetate] to give the desired product (1.02g), as a yellow oil. δ H (CDCl₃) 8.51 (1H,

d, \downarrow 4.9Hz), 7.21 (1H, br s), 6.92 (3H, br s), 3.86 (6H, s), 3.82 (3H, s), 3.69 (3H, br s) and 3.36 (3H, s). MS m/z 349 (M+H)⁺.

The 4-carboxy-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine hydrochloride was prepared by heating a solution of 4-cyano-N-(3,4,5-trimethoxyphenyl)-N-pyrimidineamine (2.88g, 10.49mmol) in aqueous 2N NaOH (60ml) and ethanol (10ml) at reflux for 3h. On cooling the reaction mixture was adjusted to pH 3 with 2M hydrochloric acid, and the resulting precipitate collected and dried to give the desired product (2.7g) as an orange solid m.p. 241°. MS m/z 306 (M+H)⁺.

The 4-cyano-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine was prepared by heating a solution of 2-chloro-4-cyanopyrimidine [G. Davies *et al.* J. Het. Chem. 1, 130-133, (1963)] (11.0g, 78.8mmol), 3,4,5-trimethoxy aniline (14.4g, 79.4mmol) and triethylamine (12.0ml) in ethanol at reflux for 4h. On cooling the resulting precipitate was collected and dried to give the desired product (13.70 g) as a yellow solid m.p. 165°. MS m/z 287 (M+H)⁺.

EXAMPLE 129

N-Phenyl-2-(3,4,5-trimethoxyphenylamino)pyrimidine-4-carboxamide

In a manner analogous to the preparation of the pyrimidine-4-carboxamide intermediate of Example 128, from 4-carboxy-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine hydrochloride (1.0g, 2.92mmol), isobutylchloroformate (0.4ml, 3.07mmol), aniline (0.28ml, 3.07mmol) and triethylamine (1.22ml, 8.76mmol) to give the title compound (655mg) as a yellow solid m.p. 158-159°. δ H (CDCl₃) 9.66 (1H, br s), 8.67 (1H, d, \downarrow 4.9Hz), 7.71-7.67 (2H, m), 7.58 (1H, d, \downarrow 4.9Hz), 7.40-7.34 (3H, m), 7.19-7.14 (1H, m), 6.88 (2H, s), 3.87 (6H, s) and 3.86 (3H, s). MS m/z 381 (M+H)⁺.

EXAMPLE 130

4-Phenylsulphonamido-N2-(3,4,5-trimethoxyphenyl)-2-pyrimidine-amine

A solution of 2-chloro-4-phenylsulphonamidopyrimidine (250mg, 0.93mmol) and 3,4,5-trimethoxyaniline (187mg, 1.02mmol) in ethoxyethanol (5ml) was heated at 140° for 12h. After this time the solvent was evaporated under reduced pressure and the residue

partitioned between ethyl acetate (10ml) and water (10ml). The organic layer was washed with water (2x25ml), dried (MgSO₄) and evaporated and the resulting solid subjected to column chromatography [silica-ethyl acetate] to give the title compound (187mg) as a buff solid, m.p. 222-223°. δ H (CDCl₃) 8.11 (1H, d, \downarrow 5.9Hz), 7.92 (3H, m), 7.57 (1H, m), 7.50 (3H, m), 6.80 (2H, s), 6.66 (1H, d, \downarrow 5.9Hz), 3.81 (6H, s) and 3.21 (3H, s). MS m/z 417 (M+H)⁺.

The 2-chloro-4-phenylsulphonamidopyrimidine was prepared by heating 2,4-dichloropyrimidine (1.42g, 9.5mmol), benzenesulphonamide (4.5g, 29.0mmol) and potassium carbonate (3.3g, 24.0mmol) in DMA (25ml) for 0.75h. The reaction was cooled to 5°, water (20ml) added, and adjusted to pH2.5 with 2M HCl. The resulting precipitate was collected to give the desired product (1.8g) as a light yellow solid m.p. 164°. MS m/z 270 (M+H)⁺.

EXAMPLE 131

N4-(tert-Butoxycarbonyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidine-diamine

To a solution of diphenylphosphorylazide (0.92ml, 4.3mmol) in *tert*-butyl alcohol (10ml), was added triethylamine (1.2ml, 8.5mmol) and 4-carboxy-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine (1.0g, 2.9mmol), [see Example 121] and the mixture heated at reflux for 2h. After this time the solvent was evaporated and the residue partitioned between ethyl acetate (100ml) and water (100ml). The organic layer was washed with water (100ml), dried (MgSO₄) and concentrated under reduced pressure. Column chromatography [silica 2% methanol CH₂Cl₂] gave the title compound (850mg) as a light yellow solid m.p. 124°. δ H (CDCl₃) 8.27 (1H, d, \downarrow 5.8Hz), 7.45 (1H, br s), 7.32 (1H, d, \downarrow 5.8Hz), 7.20 (1H, br s), 6.81 (2H, s), 3.85 (6H, s), 3.81 (3H, s) and 1.53 (9H, s). MS m/z 377 (M+H)⁺.

EXAMPLE 132

4-Phenylcarboxamido-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine

In a manner analogous to the preparation of the pyrimidine-4-carboxamide intermediate of Example 128, from benzoic acid (66mg, 0.54mmol), isobutylchloroformate (0.078ml, 0.6mmol) N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine (150mg, 0.54mmol) to give the title compound

(30mg) as a white solid m.p. 88-89°. δ H (CDCl₃) 8.39 (1H, d, \downarrow 5.6Hz), 8.32 (1H, br s), 7.89-7.85 (2H, m), 7.74 (1H, d, \downarrow 5.6Hz), 7.64-6.50 (3H, m), 6.99 (1H, br s), 6.85 (2H, s), 3.88 (6H, s) and 3.84 (3H, s). MS m/z 381 (M+H)⁺.

- 5 The N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine used as starting material was prepared by treating a solution of the compound of Example 131 (600mg, 1.6mmol) in methanol (10ml) with 2M hydrochloric acid (10ml), which was then heated at reflux for 2h. On cooling the methanol was evaporated and the aqueous solution adjusted to pH 12 with 2N
10 aqueous NaOH. The basic solution was extracted with CH₂Cl₂ (3x30ml), then the combined organic extracts were washed with saturated brine (1x20ml), dried (MgSO₄) and concentrated under reduced pressure to give the desired product (340mg) as a light yellow solid m.p. 89°. MS m/z 277 (M+H)⁺.

15

EXAMPLE 133

N2-(3,4-Dimethoxy-5-hydroxy)-N4-(3-methoxyphenyl)-2,4-pyrimidine-diamine

- In a manner analogous to the preparation of the compound of Example 49,
20 from the compound of Example 117 (200mg, 0.44mmol) to give the title compound (109mg) as a white solid m.p. 94-97°. δ H (d⁶ DMSO) 9.24 (1H, s), 8.86 (1H, s), 8.84 (1H, s), 7.97 (1H, d, \downarrow 6.0Hz), 7.32 (1H, d, \downarrow 9.2Hz), 7.26 (1H, s), 7.17 (1H, t, \downarrow 8.1Hz), 6.89 (2H, m), 6.54 (1H, d, \downarrow 8.3Hz), 6.17 (1H, d, \downarrow 5.7Hz), 5.73 (3H, s), 5.69 (3H, s) and 3.63 (3H, s).

25

The compound of Examples 134-137 were prepared in a similar manner to the compound of Example 1 from the starting materials shown:

EXAMPLE 134

5-Methoxy-4-phenyl-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine

- 30 from 3,4,5-trimethoxyphenylguanidine nitrate (1.43g, 5.0mmol), 3-dimethylamino-2-methoxy-1-phenyl-2-propen-1-one (1.03g, 5.0mmol) and NaOH (220mg) to give the title compound (200mg) as straw coloured crystals m.p. 168-170°. δ H (d⁶ DMSO) 9.36 (1H, br s), 8.47 (1H, s), 8.09 (2H, m), 7.48 (3H, s), 7.26 (2H, s), 3.88 (3H, s), 3.75 (6H, s) and 3.59 (3H, s). MS m/z 368 (M+H)⁺.

35

The 3-dimethylamino-2-methoxy-1-phenyl-2-propen-1-one was prepared in a similar manner to the analogous starting material of Example 61, from 2-methoxyacetophenone (4.51g, 30.0mmol) to give the desired product (5.30g) as an orange oil. MS m/z 206 (M+H)⁺.

5

EXAMPLE 135**4-Phenyl-5-thiomethyl-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

from 3,4,5-trimethoxyphenylguanidine nitrate (1.43g, 5.0mmol), 3-dimethylamino-1-phenyl-2-thiomethyl-2-propen-1-one (1.11g, 5.0mmol)
10 and NaOH (220mg, 5.5mmol) to give the title compound (79.8mg) as light yellow crystals m.p. 143°. δ H (CDCl₃) 8.52 (1H, s), 7.88-7.85 (2H, m), 7.49 (3H, m), 7.16 (1H, br s), 7.02 (2H, s), 3.86 (6H, s), 3.82 (3H, s) and 2.26 (3H, s). MS m/z 384 (M+H)⁺.

The 3-dimethylamino-1-phenyl-2-thiomethyl-2-propen-1-one was prepared
15 in a similar manner to the analogous starting material of Example 61, from 2-thiomethylacetophenone (4.28g, 25.7mmol) to give the desired product (3.41g) as an orange oil. δ H (CDCl₃) 7.49-7.47 (2H, m), 7.46 (1H, s), 7.41-7.34 (3H, s), 3.27 (6H, s) and 2.12 (3H, s) MS m/z 222 (M+H)⁺.

20 **EXAMPLE 136****5-Nitro-4-phenyl-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

from 3,4,5-trimethoxyphenylguanidine nitrate (1.14g, 4.0mmol), 3-dimethylamino-2-nitro-1-phenyl-2-propen-1-one (0.88g, 4.0mmol) and NaOH (176mg, 4.4mmol) to give the title compound (1.33g) as an orange
25 powder m.p. 188-189°. δ H (d⁶ DMSO) 10.60 (1H, br s), 9.16 (1H, s), 7.61 (2H, m), 7.56-7.46 (3H, m), 7.25 (2H, br s), 3.74 (6H, s) and 3.63 (3H, s). MS m/z 383 (M+H)⁺.

The 3-dimethylamino-2-nitro-1-phenyl-2-propen-1-one was prepared in a similar manner to the analogous starting material of Example 61 from
30 benzoylnitromethane (4.12g, 25.0mmol) to give the desired product (1.27g) as yellow crystals m.p. 103-105°. MS m/z 243 [(M+Na)⁺, 70%] 221 [(M+H)⁺, 100%].

EXAMPLE 137**5-Ethoxycarbonyl-4-phenyl-N-(3,4,5-trimethoxyphenyl)-2-pyrimidine-amine**

- from 3,4,5-trimethoxyphenylguanidine nitrate (2.86g, 10.0mmol), 3-
5 dimethylamino-2-ethoxycarbonyl-1-phenyl-2-propen-1-one (2.47g,
10.0mmol) and NaOH (440mg, 11.0mmol) to give the title compound
(3.10g) as straw coloured crystals m.p. 160-161°. δ H (d^6 DMSO) 10.12
(1H, br s), 8.87 (1H, s), 7.59 (2H, m), 7.50-7.45 (3H, m), 7.28 (2H, s), 4.11
(2H, q, \downarrow 7.1Hz), 3.74 (6H, s), 3.61 (3H, s) and 1.06 (3H, t, \downarrow 7.1Hz).
10 The 3-dimethylamino-2-ethoxycarbonyl-1-phenyl-2-propen-1-one was
prepared in a similar manner to the analogous starting material of
Example 61, from ethylbenzoylacetate (5.77g, 30.0mmol) to give the
desired product (6.01g) as a yellow solid m.p. 64-65°. MS m/z 248
(M+H)⁺.

15

EXAMPLE 138**5-Amino-4-phenyl-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

- In a manner analogous to the preparation of the compound of Example 38,
from the compound of Example 136 (500mg, 1.32mmol) to give the title
20 compound (260mg) as a yellow solid m.p. 149-150°. δ H (CDCl₃) 8.09
(1H, s), 7.87-7.84 (2H, m), 7.51-7.47 (3H, m), 7.00 (2H, s), 6.87 (1H, br s),
3.86 (6H, s), 3.81 (3H, s) and 3.52 (2H, br s). MS m/z 353 (M+H)⁺.

EXAMPLE 139**4-Phenylsulphanyl-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine**

- In a manner analogous to the preparation of the compound of Example 89,
from 2-chloro-4-phenyl-sulphanylpuridine (1.41g, 6.35mmol) and 3,4,5-
trimethoxyphenylaniline (1.16g, 6.35mmol) to give the title compound
(1.43g) as an off-white solid m.p. 110°. δ H (CDCl₃) 8.04 (1H, d, \downarrow 5.4Hz),
30 7.59 (2H,m), 7.47 (3H, m), 7.22 (1H, br s), 6.88 (2H, s), 6.18 (1H, d, \downarrow
5.4Hz), 3.83 (6H, s) and 3.81 (3H, s). MS m/z 370 (M+H)⁺.

- The 2-chloro-4-phenylsulphanylpuridine was prepared by treating a
solution of thiophenol (4.7ml, 45.4mmol) in THF (30ml) at 0°, with a 1.0M
solution of sodium bis(trimethylsilyl)amide in THF (45.4ml) and the mixture
35 stirred for 0.5h. After a thick white precipitate had formed, 2,4-
dichloropyrimidine (6.83g, 45.4mmol) and DMA (70ml) were added and

the mixture heated at 120° for 2h. The reaction was then concentrated under reduced pressure and the residue partitioned between water and ethyl acetate. The aqueous layer was further extracted with ethyl acetate and the combined organic layers were dried (MgSO₄) and evaporated.

- 5 The residue was subjected to column chromatography (silica 20% ethyl acetate-hexane) to give the desired product (2.66g) as a white solid. δ H 8.17 (1H, d, \downarrow 5.5Hz), 7.59 (2H, m), 7.53 (3H, m) and 6.62 (1H, d, \downarrow 5.5Hz). MS m/z 223 (M+H)⁺.

10 **EXAMPLE 140**

N-(4-(2-(3,4,5-Trimethoxyphenylamino)pyrimidin-4-yl)-2-amino-acetamide

- A solution of N-(4-(2-(3,4,5-trimethoxyphenylamino)pyrimidin-4-yl)phenyl)-2-N-(9-fluorenylmethoxycarbonyl)aminoacetamide (210mg, 0.33mmol) and piperidine (0.5ml, 5mmol) in DMF (2.5ml) was stirred at room temperature for 0.5h. The solvent was evaporated and the residue triturated with hot ethyl acetate to give the title compound (25mg) as an off-white solid m.p. 213°. δ H (d⁶ DMSO) 9.47 (1H, s), 8.49 (1H, d, \downarrow 5.4Hz), 8.11 (2H, d, \downarrow 8.3Hz), 7.75 (2H, d, \downarrow 8.2Hz), 7.29 (3H, m), 3.80 (6H, s) and 3.61 (3H, s). MS m/z 410 (M+H)⁺.
- 15
20

- The starting material for the above process was obtained by treating a solution of the compound of Example 39 (250mg, 0.71mmol) in CH₂Cl₂ (10ml) with N-FMOCglycyl chloride (337mg, 1.07mmol) and 5% aqueous Na₂CO₃ solution (8ml). After the mixture had been stirred for 1h at room temperature, the aqueous phase was discarded and the organic layer dried (MgSO₄) and evaporated. The residue was subjected to column chromatography [silica 10% methanol-CH₂Cl₂] to give the desired product (235mg) as an off-white solid. m.p. 102° (decomp.). δ H (d⁶ DMSO) 10.23 (1H, s), 9.48 (1H, s), 8.48 (1H, d, \downarrow 5.1Hz), 8.15 (2H, d, \downarrow 8.3Hz), 7.88 (2H, d, \downarrow 7.3Hz), 7.73 (4H, m), 7.64 (1H, m), 7.43-7.30 (7H, m), 4.34-4.20 (3H, m), 3.79 (8H, m) and 3.62 (3H, s).
- 25
30

EXAMPLE 141

6-Methyl-4-phenoxy-N-(3,4,5-trimethoxyphenyl)pyrimidine-2-amine

- 35 4-Chloro-6-methyl-N-(3,4,5-trimethoxyphenyl)pyrimidine-2-amine (0.30g, 0.97mmol) was added to a solution of sodium phenoxide (0.97mmol)

[prepared from phenol (0.091g, 0.97mmol) and NaH (60% in mineral oil; 0.039g, 0.97mmol)], and heated to 80° for 17h with stirring. Water was added and the mixture was extracted with CH₂CH₂ (2x30ml), the organic phases were dried (MgSO₄) and evaporated to a yellow oil. The residue was subjected to chromatography (silica, 2% methanol/CH₂CH₂) to give the title compound (0.046g). A portion was recrystallised from methanol to give an off white crystalline solid m.p. 175°. δ_H (CDCl₃) 2.38 (3H, s), 3.64 (6H, s), 3.77 (3H, s), 6.19 (1H, s), 6.78 (2H, s), 6.95 (1H, s), 7.15 (2H, d, J 7.5Hz), 7.23 (1H, m) and 7.39 (2H, t, J 7.5Hz).

10

The amine starting material for this reaction was prepared from the following:

4-Chloro-6-methyl-N-(3,4,5-trimethoxyphenyl)pyrimidine-2-amine (b)
6-Methyl-2-(3,4,5-trimethoxyphenylamino)-4(3H)pyrimidinone (1.0g, 3.63mmol) was dissolved in phosphoryl chloride and heated to 100° for 2 h. The mixture was cooled to room temperature and evaporated *in vacuo* to a viscous oil. The oil was dissolved in CH₂CH₂ (50ml) and washed twice with saturated aqueous NaHCO₃, the organic phases dried (MgSO₄) and concentrated *in vacuo*. The residue was subjected to column chromatography (silica, 2% methanol/CH₂CH₂) to afford the title compound as an off white solid (1.03g). A small amount was recrystallised from ethyl acetate to give fine off-white needles. m.p. 195-196°. δ_H (CDCl₃) 2.40 (3H, s), 3.82 (3H, s), 3.88 (6H, s), 6.64 (1H, s), 6.95 (2H, s), and 7.07 (1H, br s).

25

The pyrimidinone starting material for this reaction was prepared as follows:

6-Methyl-2-(3,4,5-trimethoxyphenylamino)-4-(3H)pyrimidinone
To a solution of sodium methoxide (0.96g, 1.79mmol) in methanol was added ethyl-3-oxo-butanoate (0.23g, 1.79mmol) and 3,4,5-trimethoxyphenylguanidinium nitrate (0.5g, 1.79mmol). The mixture was stirred at reflux for 17h. The reaction mixture was cooled to room temperature and evaporated *in vacuo* to give a dark grey solid. Recrystallisation from ethanol gave the title compound (0.25g, 49%) as a light grey solid, m.p. 228-230° (dec). δ_H (d⁶DMSO) 2.12 (3H, s), 3.61 (3H, s), 3.74 (6H, s), 5.69 (1H, s), 7.02 (2H, s), 8.70 (1H, br s) and 10.47 (1H, br s).

35

BIOLOGICAL ACTIVITY

The following assays were used to demonstrate the activity and selectivity of compounds according to the invention:

5

p56^{lck} kinase assay

The tyrosine kinase activity of p56^{lck} was determined using a RR-src peptide (RRLIEDNEYTARG) and [γ -³³P]ATP as substrates. Quantitation of the ³³P-phosphorylated peptide formed by the action of p56^{lck} was achieved using an adaption of the method of Geissler *et al* (J. Biol. Chem. (1990) 265, 22255-22261).

10

All assays were performed in 20mM HEPES pH 7.5 containing 10mM MgCl₂, 10mM MnCl₂, 0.05% Brij, 1 μ M ATP (0.5 μ Ci[γ -³³P]ATP) and 0.8mg/ml RR-src. Inhibitors in dimethylsulphoxide (DMSO) were added such that the final concentration of DMSO did not exceed 1%, and enzyme [human p56^{lck}] such that the consumption of ATP was less than 10%. After incubation at 30°C for 15min, the reaction was terminated by the addition of one-third volume of stop reagent (0.25mM EDTA and 33mM ATP in dH₂O). A 15 μ l aliquot was removed, spotted onto a P-30 filtermat (Wallac, Milton Keynes, UK), and washed sequentially with 1% acetic acid and dH₂O to remove ATP. The bound ³³P-RR-src was quantitated by scintillation counting of the filtermat in a Betaplate scintillation counter (Wallac, Milton Keynes, UK) after addition of Meltilex scintillant (Wallac, Milton Keynes, UK). The dpm obtained, being directly proportional to the amount of ³³P-RR-src produced by p56^{lck}, were used to determine the IC₅₀ for each compound.

15

20

25

p59^{fyn} kinase assay

Compounds of the invention were assayed for p59^{fyn} inhibitory activity in a similar manner to the p56^{lck} assay, using human p59^{fyn}.

30

EGFr kinase assay

The tyrosine kinase activity of the EGF receptor (EGFr) was determined using a similar methodology to the p56^{lck} kinase assay, except that the RR-src peptide was replaced by a peptide substrate for EGFr obtained

35

from Amersham International plc (Little Chalfont, UK) and used at the manufacturers recommended concentration. IC₅₀ values for each test inhibitor were determined as described previously in the p56^{lck} assay.

5 **ZAP-70 kinase assay**

The tyrosine kinase activity of ZAP-70 was determined using a capture assay based on that employed above for p56^{lck}. The RR-src peptide was replaced with polyGlu-Tyr (Sigma; Poole, UK) at a final concentration of 17 µg/ml. After addition of the stopped reaction to the filtermat, trichloroacetic acid 10% (w/v) was employed as the wash reagent instead of acetic acid and a final wash in absolute ethanol was also performed before scintillation counting. IC₅₀ values for each test inhibitor were determined as described above in the p56^{lck} assay.

10

15 **Csk kinase assay**

Compounds of the invention were assayed for csk kinase inhibitory activity in a similar manner to the ZAP-70 assay using human csk kinase.

Protein kinase C assay

Inhibitor activity against protein kinase C (PKC) was determined using PKC obtained from Sigma Chemical Company (Poole, UK) and a commercially available assay system (Amersham International plc, Little Chalfont, UK). Briefly, PKC catalyses the transfer of the γ-phosphate (32P) of ATP to the threonine group on a peptide specific for PKC. Phosphorylated peptide is bound to phosphocellulose paper, subsequently quantified by scintillation counting and IC₅₀ values determined as before.

20

25

In the above assays, compounds according to the invention, including the compounds of the Examples inhibit the protein kinase p56^{lck}, p59^{fyn}, ZAP-70 or protein kinase C at concentrations at which they have little or no effect on the remaining kinases, including EGFr kinase and Csk kinase. Thus, for example the compounds of Example 96 and 103 have IC₅₀ values in the p56^{lck} assay of 215nM and 40nM respectively. The compounds are also active against p59^{fyn} but have IC₅₀ values at least 10x greater against the remaining kinases described above. In another example the compounds of Example 64 and 66 have IC₅₀ values in the


30

35

ZAP-70 assay of 124nM and 68nM respectively. The compounds are also active in the protein kinase C assay, but have IC₅₀ values at least 10x greater against the remaining kinases. In a further example, the compounds of Examples 48 and 81 have IC₅₀ values in the protein kinase C assay of 22nM and 92nM respectively, but have IC₅₀ values at least 10x greater against the remaining kinases.

CLAIMS

1. A compound of formula (1):



(1)

wherein R¹ is a hydrogen or halogen atom or an optionally substituted straight or branched chain alkyl, alkenyl or alkynyl group or a group selected from hydroxyl (-OH), substituted hydroxyl, thiol (-SH), substituted thiol, amino (-NH₂), or substituted amino;

R² and R³, which may be the same or different, is each an optionally substituted straight or branched chain alkyl, alkenyl or alkynyl group;

R⁴ is a hydrogen atom or a straight or branched chain alkyl group;

R⁵ is a hydrogen atom or an optionally substituted straight or branched chain alkyl, alkenyl or alkynyl group;

R⁶ is a hydrogen or halogen atom or an amino (-NH₂), substituted amino, nitro, carboxyl (-CO₂H) or esterified carboxyl group or a group -X¹-R^{6a} where X¹ is a direct bond or a linker atom or group and R^{6a} is an optionally substituted straight or branched chain alkyl, alkenyl or alkynyl group;

X is a direct bond or a linker atom or group;

R⁷ is an optionally substituted aliphatic, cycloaliphatic, heteroaliphatic, heterocycloaliphatic, aromatic or heteroaromatic group;

and the salts, solvates, hydrates and N-oxides thereof.

2. A compound according to Claim 1 wherein R⁴ is a hydrogen atom.

3. A compound according to Claim 1 or Claim 2 wherein R⁵ and R⁶ is each a hydrogen atom.
- 5 4. A compound according to any one of Claims 1 to 3 wherein R¹ is an optionally substituted methoxy group and R² and R³ is each an optionally substituted methyl or ethyl group.
- 10 5. A compound according to Claim 4 wherein R¹ is a methoxy group and R² and R³ is each a methyl group.
- 15 6. A compound according to any one of the preceding Claims wherein X is a direct bond, a sulphur atom or a -NH- group.
7. A compound according to any one of Claims 1 to 6 wherein R⁷ is an optionally substituted aromatic or heteroaromatic group.
- 20 8. A compound according to Claim 7 wherein R⁷ is an optionally substituted phenyl, 2-pyridyl, 3-pyridyl or 4-pyridyl group.
- 25 9. A compound which is
N,N'-Bis(3,4,5-Trimethoxyphenyl)-2,4-pyrimidinediamine;
4-(2-(2-Dimethylaminoethylamino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine;
4-(2-(4-Hydroxybutylamino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine;
4-(3-(3-Aminopropoxy)phenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine;
N4-(3,4-Dimethoxyphenyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine;
30 4-(4-(2-Hydroxyethoxy)phenyl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine;
4-(2-(3-(Morpholino)propylamino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine;
35 4-(2-(2-(1-Methylpyrrolidin-2-yl)ethylamino)pyridin-5-yl)-N-(3,4,5-trimethoxyphenyl)-2-pyrimidineamine;

N4-(4-(Ethoxycarbonylmethyl)phenyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine;


N4-(4-Hydroxyphenyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidine-diamine and

5 N4-(4-(3-Aminopropoxy)phenyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine;

and the salts, solvates, hydrates and N-oxides thereof.

10. A pharmaceutical composition comprising a compound of formula 1:

10



(1)

wherein R¹ is a hydrogen or halogen atom or an optionally substituted straight or branched chain alkyl, alkenyl or alkynyl group or a group selected from hydroxyl (-OH), substituted hydroxyl, thiol (-SH), substituted thiol, amino (-NH₂), or substituted amino;

15

R² and R³, which may be the same or different, is each an optionally substituted straight or branched chain alkyl, alkenyl or alkynyl group;

R⁴ is a hydrogen atom or a straight or branched chain alkyl group;

20

R⁵ is a hydrogen atom or an optionally substituted straight or branched chain alkyl, alkenyl or alkynyl group;

R⁶ is a hydrogen or halogen atom or an amino (-NH₂), substituted amino, nitro, carboxyl (-CO₂H) or esterified carboxyl group or a group -X¹-R^{6a} where X¹ is a direct bond or a linker atom or group and R^{6a}

25

is an optionally substituted straight or branched chain alkyl, alkenyl or alkynyl group;

X is a direct bond or a linker atom or group;

- R⁷ is an optionally substituted aliphatic, cycloaliphatic, heteroaliphatic, heterocycloaliphatic, aromatic or heteroaromatic group;
and the salts, solvates, hydrates and N-oxides thereof;
- 5 together with one or more pharmaceutically acceptable carriers, excipients or diluents.

INTERNATIONAL SEARCH REPORT

Internat'l Application No
PCT/GB 96/02854

A. CLASSIFICATION OF SUBJECT MATTER IPC 6 C07D239/42 C07D239/48 C07D401/04 C07D417/04 C07D409/04 A61K31/505		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 6 C07D		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	FR,A,2 545 356 (RIPHARM) 9 November 1984 see page 1 - page 7 ---	1,10
A	EP,A,0 295 210 (CIBA-GEIGY) 14 December 1988 see claims; examples 13,29,41,49,6 -----	1,10
<input type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents :		
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>*A* document defining the general state of the art which is not considered to be of particular relevance</p> <p>*E* earlier document but published on or after the international filing date</p> <p>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>*O* document referring to an oral disclosure, use, exhibition or other means</p> <p>*P* document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>*&* document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search <div style="text-align: center; font-weight: bold;">4 February 1997</div>		Date of mailing of the international search report <div style="text-align: center; font-weight: bold;">12.02.97</div>
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl, Fax (+ 31-70) 340-3016		Authorized officer <div style="text-align: center; font-weight: bold;">Francois, J</div>

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/GB 96/02854

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
FR-A-2545356	09-11-84	DE-A- 3414629	22-11-84
EP-A-295210	14-12-88	AU-B- 615912	17-10-91
		AU-A- 1760688	15-12-88
		BG-A- 48205	14-12-90
		BG-A- 49923	16-03-92
		BG-A- 49939	16-03-92
		CA-A- 1305483	21-07-92
		CN-A- 1032338	12-04-89
		CZ-A- 8803926	11-09-96
		DE-A- 3882821	09-09-93
		ES-T- 2058332	01-11-94
		HU-B- 208961	28-02-94
		IE-B- 62436	08-02-95
		JP-A- 63316774	26-12-88
		PT-B- 87707	31-01-95